

INTRODUCTION
This Surficial Geology Map of NTS 94-09 (Canadian Geoscience Map 123) is the product of collaboration between the Geological Survey of Canada, the British Columbia Ministry of Energy, Mines and Natural Gas as part of the Geoscience Mapping for Energy and Minerals Program (GEM) - Energy Yukon Basins Project.

APPROACH TO SURFICIAL GEOLOGY MAPPING
The surficial and bedrock-based geomorphic features have led to a better understanding of the regional distribution of surficial deposits, permafrost, landslides and other geomorphic processes in the NTS 94-09 map area (Huntley and Hickin, 2010; Huntley et al., 2011a-b).

INFERRED GEOLOGICAL HISTORY
The distinctive landscape of NTS 94-09 largely a product of underlying bedrock and geological structures, with ornamentation by the Upper Wisconsinan Laurentide Ice Sheet. The Eshbo Plateau is underlain by conglomerate, sandstone and carbonaceous shale of the Lake Wisconsinan Formation.

ACKNOWLEDGMENTS
Canadian Geoscience Map 123 is an output of the Geo-Mapping for Energy and Minerals Yukon Basins Project managed by Carl Ozyer and Amy Law (GSC-Canada). The assets (Data, Maps, Images, etc.) were provided by the Geological Survey of Canada (GSC-Canada).

Abstract
Canadian Geoscience Map 123 depicts the surficial geology over some 700 km² covered by the Trail Lake map sheet (NTS 94-09), in northeastern British Columbia. Here, the Eshbo Plateau lies within the Pelletier River watershed and is drained by east-flowing Gole Creek, and northward by Dilly and Yesashalle creeks.

Résumé
La Carte géoscientifique du Canada 123 illustre la géologie des matériaux superficiels d'un territoire d'environ 700 km² couvert par le feuillet cartographique de Trail Lake (SNRC 94-09), dans le nord-est de la Colombie-Britannique. Ici, le plateau d'Eshbo se situe dans le bassin versant de la rivière Pelletier et est drainé par le ruisseau Gole vers l'est ainsi que par les ruisseaux Dilly et Yesashalle vers le nord.

National Topographic System reference and index to adjoining published Geological Survey of Canada maps

Cover Illustration
Rotational landslide triggered by the Kwigwina River incising the Eshbo Plateau in northeast British Columbia, view west. Photograph by D.H. Huntley, 2013-101

Canadian Geoscience Map 123 is an output of the Geo-Mapping for Energy and Minerals Yukon Basins Project managed by Carl Ozyer and Amy Law (GSC-Canada). The assets (Data, Maps, Images, etc.) were provided by the Geological Survey of Canada (GSC-Canada).

Authors: D.H. Huntley, A.S. Hickin, W. Chow, and M. Mirrezaei
Geomatics by D.H. Huntley, W. Chow, and M. Mirrezaei
Cartography by W. Chow

Geological Survey of Canada
Initiative of the Geological Survey of Canada, conducted under the auspices of the Yukon Basins Project as part of Natural Resources Canada's Geomapping for Energy and Minerals (GEM) program

Map projection Universal Transverse Mercator, zone 10, North American Datum 1983
Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications
Elevations in feet above mean sea level
Magnetic declination 2013, 2019'E, decreasing 22' annually.

The Geological Survey of Canada welcomes corrections or additional information from users.
This publication is available for free download through GEOCAN (http://geocan.ess.nrcan.gc.ca/)

Recommended citation
Huntley, D. H., Hickin, A. S., Chow, W., and Mirrezaei, M., 2013. Surficial geology, Trail Lake, British Columbia, Geological Survey of Canada, Canadian Geoscience Map 123 (preliminary), scale 1:50 000, doi:10.4095/262400

ess.nrcan.gc.ca
Natural Resources Canada / Ressources naturelles du Canada

CANADIAN GEOSCIENCE MAP 123
SURFICIAL GEOLOGY
TRAIL LAKE
British Columbia
1:50 000


1:50 000
Scale bar showing 0 to 4 km

1:50 000
Scale bar showing 0 to 4 km

1:50 000
Scale bar showing 0 to 4 km

1:50 000
Scale bar showing 0 to 4 km

1:50 000
Scale bar showing 0 to 4 km



Bednarski, J.M., 2003a. Batulama Lake, Northwest Territories - Yukon Territory - British Columbia (NTS 9504). Geological Survey of Canada, Open File 4502, scale 1:50 000.
Bednarski, J.M., 2003b. Surficial geology of Fort Liard, Northwest Territories - British Columbia, Geological Survey of Canada, Open File 1790, scale 1:50 000.
Bednarski, J.M., 2003c. Surficial geology of Lake Bovin, Northwest Territories - British Columbia, Geological Survey of Canada, Open File 1791, scale 1:50 000.
Bednarski, J.M., 2003d. Surficial geology of Colville Lake, Northwest Territories - British Columbia, Geological Survey of Canada, Open File 1794, scale 1:50 000.
Bednarski, J.M., 2003e. Surficial geology of Etah Creek, British Columbia, Geological Survey of Canada, Open File 4844, scale 1:50 000.
Clemens, C., Kowal, R., Huntley, D. and Dalziel, R., 2004. Ecosystem units of the Saitheen area, Sitka Forest Products (Fort Nelson Report), 39 pages and appendices.
Dobbie, C., Proulx, A., Boisvert, E., Butler, G., Davernop, P., Everett, D., Huntley, D., Inglis, E., Kerr, D., Moore, A., Paradis, S.J., Parent, M., Smith, R., St-Onge, D., and Weatherman, A., 2012. Science Language for an Integrated Geological Survey of Canada Data Model for Surficial Maps Version 1.1. Technical Report of the Geological Survey of Canada Surficial Legend Review Committee, Geological Survey of Canada, Open File 7003, 27 pages.
Demchuk, T., 2010. Surficial geology of the Komie Creek area (NTS 04P05). British Columbia Ministry of Energy, Mines and Petroleum Resources, Geological Survey of Canada Open File 6668, scale 1:50 000.
Huntley, D.H. and Hickin, A.S., 2010. Surficial deposits, landforms, glacial history and potential for glacial aggregate and fine sand: Muwashish Lake Map Area (NTS 94-0), British Columbia, Geological Survey of Canada, Open File 6430, 17 pages.
Huntley, D.H. and Chow, W., 2014. Surficial geology, geomorphology, granular resource evaluation and geohazard assessment for the Muwashish Lake map area (NTS 94-0), northeastern British Columbia, Geological Survey of Canada, Open File 6982, 20 pages.
Huntley, D.H., Hickin, A.S. and Fren, F., 2010. Provisional surficial geology, glacial history and geomorphological reconstructions of the Tsai River (NTS 94-0) and Muwashish Lake map areas (NTS 94-0), British Columbia, Geoscience Report 2011, 102 Ministry of Energy, pages 37-55.
Huntley, D.H. and Sobel, C.F., 2010. Application of the GEM surficial geology data model to resource evaluation and geohazard assessment for the Muwashish Lake map area (NTS 94-0), British Columbia, Geological Survey of Canada, Open File 6553, 22 pages.
Stott, D.F. and Taylor, C.E., 1968. Geology of Muwashish Lake, Geological Survey of Canada, Paper 1-1968, scale 1:250 000.
Tomlinson, M. and Levason, V.M., 2008. Quaternary stratigraphy of the Proglacial River, northeastern British Columbia, Canadian Journal of Earth Sciences, Vol. 45, pages 945-975.

Late Pleistocene earth materials and landforms
Glaciolacustrine deposits
Silt and clay with subordinate sand, gravel and diamicton, massive or rhythmically interbedded, slump structures and drapings locally present; poor to moderately drained, generally greater than 2 m thick, kettle lakes and irregular topography underlain by bedrock, till and outwash; transported by and deposited from sediment laden meltwater; subaqueous gravity flows and normal melting of ice in proglacial lakes; were sporadically discontinuous permafrost is, or was present, glaciolacustrine sediments may be subject to thermokarst processes; slopes less than 5° are potentially unstable and prone to landslides and debris flows.

Glaciolacustrine blanket: silt and clay with subordinate sand, gravel and diamicton, massive or rhythmically interbedded, slump structures and drapings locally present; poor to moderately drained, generally greater than 2 m thick, kettle lakes and irregular topography underlain by bedrock, till and outwash; transported by and deposited from sediment laden meltwater; subaqueous gravity flows and normal melting of ice in proglacial lakes; were sporadically discontinuous permafrost is, or was present, glaciolacustrine sediments may be subject to thermokarst processes; slopes less than 5° are potentially unstable and prone to landslides and debris flows.

Kames and hummocky outwash: boulders, cobbles, pebble-gravel, sand, silt and diamicton, generally massive to stratified, some slump structures; moderately to well-drained, greater than 2 m thick, terrace scarpings range from 100 m to 4 km in length; in contact with, and overlying till units, outwash and glaciolacustrine sediments, deposited by rivers and granular aggregate with glacial ice; potential source of groundwater and granular aggregate when material is gravel rich.

Esker terraces: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamicton, generally massive to stratified, some slump structures; moderately to well-drained; greater than 2 m thick; range from 100 m to 8 km in length; in contact with, and overlying till units, outwash and glaciolacustrine sediments, deposited by subglacial meltwater in contact with glacial ice; potential source of groundwater and granular aggregate when material is gravel rich.

Outwash terraces: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamicton, generally massive to stratified, some slump structures; moderately to well-drained; greater than 2 m thick; terrace scarpings range from 100 m to 4 km in length; in contact with, and overlying till units, outwash and glaciolacustrine sediments, deposited by meltwater confined to proglacial channels and spillways; potential source of groundwater and granular aggregate when material is gravel rich.

Till deposits
Till blanket: sand, silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; transported and deposited by the Laurentide Ice Sheet directly through lodgement, basal meltout, glaciogenic deformation of sediment beneath active, warm-based ice and in situ melting from stagnant cold-based ice; stable terrain, generally suitable for infrastructure placement.

Hummocky till: sand and silt-rich diamictons; massive to stratified, matrix- and clast-supported; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; drapes till and other glacial deposits; deposited by silt melting from stagnant cold-based ice and modified by meltwater; evidence for ice collapse includes slump structures, kettle lakes and irregular topography; potential source of aggregate when material is gravel rich; generally suitable for infrastructure placement.

Moraine ridges: sand, silt and clay-rich diamictons; massive, matrix-supported; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; minor moraines less than 1 km long and 5 m high; major moraines up to 12.5 km in length and 10 m high; ridges drapage bedrock and older glacial deposits; minor moraines include crosswise ridges and small recessional push moraines; major ridges features are large recessional end moraines and ice-trust ridges; generally suitable for infrastructure placement.

Streamlined till: silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; drumlins and fluted till ridges typically under 1 km long but can exceed 8 km in length; generally less than 50 m wide and 20 m high; formed beneath the Laurentide Ice Sheet directly through lodgement, basal meltout, glaciogenic deformation of sediment beneath rapidly-flowing warm-based ice; generally suitable for infrastructure placement.

Geological boundary (Confidence: approximate)
Major moraine ridge (unspecified)
Other moraine ridge (unspecified)
Esker ridge (sense: unknown or unspecified)
Drumlin ridge
Major meltwater channel scarp
Minor meltwater channel central axis (unspecified; sense: known)
Station location (ground observation or stratigraphic section)

Recommended citation
Huntley, D. H., Hickin, A. S., Chow, W., and Mirrezaei, M., 2013. Surficial geology, Trail Lake, British Columbia, Geological Survey of Canada, Canadian Geoscience Map 123 (preliminary), scale 1:50 000, doi:10.4095/262400

Preliminary publications in this series have not been scientifically edited.

CANADIAN GEOSCIENCE MAP 123
SURFICIAL GEOLOGY
TRAIL LAKE
British Columbia