FORMALLY NAMED TECTONIC SETTING

Unit 2 Label

Unit 1 Label

ASSEMBLAGE AGE, PERIOD: Series (Stage, Absolute age range)

ASSEMBLAGE NAME: *assemblage description. in the following form: major rock types; minor rock types; setting (**constituent "...group", formation names indicated in bold text**); assemblages are subjective groupings of stratigraphic or metamorphic rock units that allow correlation and portrayal of lithological continuity on a regional scale*
*2. Separately distinguished and described components of the assemblage are indicated by units (i.e. † 2, 2 ...) each unit description, in the following form: major rock types; and component features; minor rock types and features; rare rock types and features; settings (**constituent "...group", formation, "...member", "...suite", "...bed" names indicated in bold text**)*
3. where indicated, some units within an assemblage are distinguished by their own colour to add detail and clarify to the map

INFORMALLY named tectonic setting

PLUTONIC ASSEMBLAGE AGE, PERIOD: Series (Stage, Absolute age range)

PLUTONIC ASSEMBLAGE NAME: *plutonic assemblage description. in the following form: major rock types; minor rock types; setting (**constituent "...suite", "...swarm", "...intrusion", "...intrusive complex" names indicated in bold text**)*
plutonic assemblages are subjective groupings of intrusive rock units that allow correlation and portrayal of lithological continuity on a regional scale
2. Separately distinguished and described components of the assemblage are indicated by units (i.e. † 2, 2 ...) each unit description, in the following form: major rock types
*and component features; minor rock types and features; rare rock types and features; settings (**constituent "...suite", "...swarm", "...intrusion", "...intrusive complex" names indicated in bold text**)*
3. where indicated, some units within an assemblage are distinguished by their own colour to add detail and clarify to the map

For clarity at this map scale, not all units in the legend are indicated separately on the map. For example, units of the Scoresby Bay assemblage, labelled Cm-ob1, Cm-ob2 and Cm-ob3 are amalgamated under the colour of their parent (Cm-ob).

Brackets indicate grouping of assemblages and component units by type of tectonic setting and colour coded as follows:

- gold: overlap assemblages (Corcoran only)
- red: plutonic etc. stitching plutons
- brown: fold/thrust belt-derived clastic wedge
- light green: rifts grading to shallow or oceanic)
- purple: slope and deep water settings
- light blue: shelf platform
- grey: intracratonic basin
- orange: rift-related
- green: large igneous province (LIP)
- dark blue: orogen, craton



National Topographic System reference and index to adjoining published Geological Survey of Canada maps

Abstract

This map and the related geodatabase illustrate the bedrock geology of central and eastern Prince of Wales Island, Somerset Island, and northern Boothia Peninsula. Major features of the area include Archean and Paleoproterozoic granitoid and metasedimentary rocks of Boothia Uplift overlain unconformably by Paleoproterozoic and Mesoproterozoic sediments, which are, in turn, unconformably overlain by Cambrian to Devonian strata of Arctic Platform. The higher Silurian and Lower Devonian interval provides a multiphase depositional record of Boothia Uplift. Younger events are recorded by Cretaceous kimberlite diatremes and outliers of Paleogene strata.

Résumé

Cette carte et la géodatabase qui s'y rapporte documentent la géologie du substratum rocheux dans le centre et l'est de l'île Prince of Wales, l'île Somerset et le nord de la presqu'île de Boothia. Les principales entités géologiques de la région comprennent des roches granitoïdes et métasédimentaires de l'Archéen et du Paléoprotérozoïque du soulèvement de Boothia, surmontées en discordance par des sédiments paléoprotérozoïques et mésoprotérozoïques qui, à leur tour, sont recouverts en discordance par des strates d'âge cambrien à dévonien de la Plate-forme de l'Arctique. L'intervalle du Silurien supérieur et du Dévonien inférieur témoigne de l'histoire sédimentaire polyphasée du soulèvement de Boothia. Des diatrèmes de kimberlite du Crétacé et des lambeaux d'érosion de strates du Paléogène rendent compte des événements les plus récents.

Cover illustration
Paleoproterozoic basement including metasediments and marble, Bellot Strait and southern Somerset Island, Nunavut. Photograph by Tom Frisch, 2013-073

Catalogue No. M183-1/76-2013E-PDF
ISBN 978-1-100-22353-7
doi:10.4095/292819

© Her Majesty the Queen in Right of Canada 2013

Natural Resources Canada / Ressources naturelles du Canada

CANADIAN GEOSCIENCE MAP 76

GEOLOGY

TECTONIC ASSEMBLAGE MAP OF PEEL SOUND

Prince of Wales and Somerset islands, Nunavut

1:500 000



Preliminary Preliminary Preliminary

PALEOGENE:

PALEOCENE AND EOCENE (Selandian to middle Eocene)

- **Te1** — **EUREKA SOUND:** sandstone, siltstone, shale; local conglomerate; minor coal, volcanogenic sandstone; mostly fluvial and deltaic; local syntectonic alluvial fan deposits.
  1. Sandstone, fine- to coarse-grained; siltstone, shale; local conglomerate, coal, volcanogenic sandstone; common plant fossils; mostly fluvial and deltaic; local syntectonic alluvial fan deposits **(Eureka Sound)**.

CRETACEOUS

EARLY AND LATE CRETACEOUS (Albian and Cenomanian; 108 to 98 Ma)

- **Kak** — **SOMERSET** *plutonic assemblage:* Kimberlite; diatreme **(Somerset kimberlite suite)**.

SILURIAN AND DEVONIAN

LUDLOW TO EARLY DEVONIAN (*Pragian*)

BOOTHIA UPLIFT

- **SDp3** — **PEEL SOUND:** sandstone, conglomerate, minor shale, siltstone, dolostone; alluvial fans, deltaic, shallow marine.
  3. Conglomerate, thick-bedded, red-weathering, carbonate clasts of granule to pebble and rare cobble grade; minor tan and grey sandstone; syntectonic alluvial fans **(Peel Sound conglomerate)**.
- **SDp2** — **2.** Sandstone, fine- to medium-grained with chert and limestone grains, yellow to green-grey and red weathering; minor conglomerate and limestone; syntectonic alluvial deposits grading to shallow marine **(Peel Sound sandstone)**.
- **SDp1** — **1.** Sandstone, fine- to medium-grained with chert and limestone grains, yellow to green-grey and red weathering; conglomerate, thick-bedded, carbonate clasts of granule to pebble and rare cobble grade; dolostone, fine-grained, laminated, desiccation cracks, molar tooth structure; minor limestone, burrowed with stromatoporoids, gastropods in the lower part; syntectonic alluvial fans grading to shallow marine, mostly restricted **(Peel Sound)**.

LUDLOW TO LOCHKOVIAN

- **SDg4** — **GOOSE FIORD:** dolostone, limestone, mudstone; minor wackestone, siltstone, sandstone; nearshore, shelf and shelf edge including reefs and foreslope, mostly unrestricted.
  4. Dolostone, medium- to thick-bedded, tan, medium-grained; limestone including lime mudstone, skeletal wackestone; packstone, fine-grained with ostracodes, crinoids, brachiopods, gastropods; silty limestone; siltstone, calcareous and dolomitic; minor sandstone, calcareous and dolomitic, very fine- to fine-grained; minor shale; nearshore to shallow offshore, unrestricted **(Drake Bay)**.

SILURIAN

LUDLOW

- **Ssi3** — **SOMERSET ISLAND:** dolostone, limestone; minor sandstone, wackestone; intertidal to supratidal.
  3. Dolostone, limestone; finely crystalline; siltstone; red, grey, green-grey, yellowish brown; thin- to medium-bedded; ripples, mud cracks, halite casts; minor sandstone, skeletal wackestone; cyclical deposition; intertidal to supratidal **(Somerset Island, Upper member)**.
- **Ssi2** — **2.** Dolostone, limestone; finely crystalline; grey, green-grey, yellowish brown; thin- to medium-bedded; ripples, mud cracks, oncolites, domal stromatolites; minor skeletal wackestone; subtidal to peritidal **(Somerset Island, Lower member)**.
- **Ssi1** — **1.** Dolostone, limestone, dolosiltite, planar laminated, grey, greenish grey, buff weathering; siltstone, mudstone; red, green, variegated; minor skeletal limestone, burrow mottled; locally common domal stromatolites, oncolites, fossil fish; rare gypsum nodules; cyclical deposition; and setting, variably shallow subtidal to intertidal **(Somerset Island)**.

- **Sdo1** — **DOURO:** Limestone; minor dolostone, shale; local bioherms; shallow to deep subtidal shelf ramp.
  1. Argillaceous limestone, limestone, dolomitic limestone; nodular, burrow-mottled, minor dolostone, calcareous shale, calcisiltite; local sponge-coral-stromatoporoid mud mounds on Somerset Island, and faveolitid-microbial biostromes in basal beds on Grinnell Peninsula; shallow to deep subtidal outer shelf ramp **(Douro)**.

- **Scs** — **CAPE STORM:** dolostone, dolomitic limestone, limestone; minor silty dolostone, siltstone, shale; arid subtidal to supratidal settings. Dolostone; grey, buff and brown weathering, very fine- to fine-grained, thin- to thick-bedded, laminated, mud-cracked, molar-tooth structures, stromatolitic, microbial; lesser dolomitic limestone, oolitic and peloidal; minor silty dolostone, siltstone, shale; mostly restricted intertidal and supratidal **(Cape Storm)**.

ORDOVICIAN TO SILURIAN

RICHMONDIAN TO LUDLOW

FRANKLINIAN-INUVIK ARCTIC PLATFORM

- **OSa2** — **ALLEN BAY:** Dolostone, limestone; local bioherms, anhydrite; arid subtidal to supratidal settings.
  2. Dolostone, laminated to thin-bedded, stromatolitic; dolostone intraclast conglomerate; anhydrite, limestone, minor siltstone; cyclical sedimentation; peritidal lagoon-cyclically grading to supratidal and hypersaline **(Cape Crauford)**.
- **OSa1** — **1.** Dolostone, medium- to thick-bedded, light grey to yellowish brown, medium-grained, in part fossiliferous; bitumen stained; lesser dolomitic limestone, burrow mottled; minor flat pebble conglomerate; local stromatoporoid bioherms; local gypsum and anhydrite (on Somerset Island); common shallowing upward cycles; semi-arid to arid subtidal to supratidal settings **(Allen Bay)**.

ORDOVICIAN

MIDDLE AND LATE ORDOVICIAN (Blackriveran to Richmondian)

- **Ocs6** — **THUMB MOUNTAIN:** limestone, mudstone, grainstone, wackestone, dolostone; subtidal and intertidal, restricted and unrestricted.
  6. Dolostone; light grey, burrow-mottled, thick-bedded; corals, stromatoporoids and cephalopods, unrestricted subtidal shelf.
- **Ocs5** — **5. Thumb Mountain, Irene Bay,** undivided.

MIDDLE ORDOVICIAN (Whiterockian)

- **Ocs1** — **BAY FIORD:** dolostone, limestone; minor anhydrite, rock salt, breccia; initially hypersaline grading to mostly restricted peritidal.
  1. Dolostone, dolomitic limestone; very fine- to fine-grained, thin- to medium-bedded, grey to grey-green weathering; limestone, pale to dark brown, massive-bedded, burrow mottled, locally bioclastic; basal discontinuous member of gypsum, anhydrite, solution breccia and, in the Parry Islands, subsurface rock salt; initially hypersaline grading up into mostly restricted peritidal **(Bay Fiord)**.

CAMBRIAN TO SILURIAN

LATE CAMBRIAN TO LUDLOW

- **CmSa** — **ARCTIC PLATFORM:** dolostone, dolomitic limestone, limestone; minor sandstone, shale, intraclast conglomerate, breccia; nearshore and intertidal to supratidal; mostly restricted **(Turner Cliffs, Ship Point, Irene Bay, Thumb Mountain, Allen Bay, Cape Storm,** undivided**)**.

CAMBRIAN AND ORDOVICIAN

LATE CAMBRIAN (Furongian) TO EARLY ORDOVICIAN (Tremadoc)

ARCTIC PLATFORM

- **CmO4** — **TURNER CLIFFS:** dolostone, limestone, intraclast conglomerate; minor shale, sandstone, gypsum; local bioherms; peritidal grading to subtidal shelf.
  4. Dolostone, very-fine to medium-grained, grey to greenish grey, variably bioturbated, arenaceous, oolitic, domal stromatolitic; locally abundant gastropods, brachiopods; minor dolomitic sandstone, dolomitic siltstone, flat pebble conglomerate, breccia; local replacement chert, local subsurface anhydrite; shallow subtidal to intertidal, mostly restricted **(Ship Point)**.
- **CmO3** — **3.** Dolostone, pale grey to buff, fine-grained, cherty, thick- to massive-bedded; resistant; shallow marine, mostly unrestricted **(Turner Cliffs, upper member)**.
- **CmO2** — **2.** Dolostone, green, grey to buff, laminated, thin- to medium-bedded, stromatolitic; sandstone, grey, white, red brown; carbonate intraclast breccia, conglomerate; nearshore to restricted shallow marine **(Turner Cliffs, lower member)**.
- **CmO1** — **1.** Dolostone, grey to buff, laminated and thin-bedded grading to massive; minor sandstone; grey, locally red brown; intraclast breccia and conglomerate; local stromatolitic units; nearshore to shallow marine **(Turner Cliffs)**; dolostone, very fine- to medium-grained, grey to greenish grey, bioturbated, arenaceous, oolitic, domal stromatolitic; gastropods, brachiopods; minor dolomitic sandstone and siltstone, intraclast conglomerate, breccia; local subsurface anhydrite; shallow subtidal to intertidal, mostly restricted **(Ship Point)**.

NEOPROTEROZOIC:

CRYOGENIAN (723 to 718 Ma)

FRANKLIN LARGE IGNEOUS PROVINCE

- **pPd** — **FRANKLIN plutonic assemblage:** Diabase, mostly high titanium tholeiite, olivine-bearing, brown to black; subophitic; plagioclase often saussuritized, olivine replaced by carbonate; local ultramafic rock; LIP intrusives, olivine tholeiite association **(Franklin and Thule swarms)**.

MESOPROTEROZOIC

ECTASIAN AND STENIAN (1270 to 1000 Ma)

BORDEN BASIN

- **mPb2** — **BORDEN:** siltstone, shale, sandstone, dolostone; syntectonic alluvial fans grading to fluvial and shallow marine.
  2. Dolostone, variably arenaceous, grey, bluish grey, buff and red weathering, thin- to thick-bedded, commonly stromatolitic, replacement chert; minor shale, red to buff; siltstone, purple; sandstone, conglomerate; shallow marine to peritidal **(Humfrey)**.

ECTASIAN (1270 Ma)

MACKENZIE LARGE IGNEOUS PROVINCE

- **mPd** — **MACKENZIE plutonic assemblage:** diabase; LIP intrusives, olivine tholeiite association. Diabase, mostly high titanium tholeiite, olivine-bearing, brown to black, subophitic; plagioclase often saussuritized, olivine replaced by carbonate **(Mackenzie swarm)**.

PALEOPROTEROZOIC TO MESOPROTEROZOIC

STATHERIAN TO ECTASIAN (1700 to 1270 Ma)

ASTON BASIN

- **pmPa** — **ASTON:** arkose, quartz sandstone; minor siltstone, dolostone; shallow marine with clastic material of local granitoid provenance. Arkose, lesser quartz sandstone, planar crossbedded, pink to red weathering, fine- to medium-grained; minor siltstone, red to purple weathering, common desiccation cracks; stromatolitic dolostone, chert in the upper part; basal breccia, conglomerate; shallow marine with clastic material of local granitoid provenance **(Aston)**.

PALEOPROTEROZOIC

OROSIRIAN TO STATHERIAN (1940 to 1700 Ma)

THELON OROGEN

- **pPm4** — **M'CLURE BAY plutonic assemblage:** anorthosite, granite, syenite; mostly syn- and post-kinematic plutonic rocks.
  4. Granite; pink, massive or porphyritic with K-feldspar phenocrysts **(M'Clure Bay suite)**.
- **pPm3** — **3.** Anorthosite; white; pale green to grey weathering; coarse-grained **(M'Clure Bay suite)**.
- **pPm2** — **2.** Sodic pyroxene syenite.
- **pPm1** — **1.** Two-pyroxene syenite.

SIDERIAN TO OROSIRIAN (2480 to 1920 Ma)

THELON OROGEN, Rae craton cover

- **pPb5** — **BOOTHIA:** paragneiss, metapelite; lesser marble; calcisilicate, granitoid gneiss; local ultramafic rocks, quartzite, meta-iron-formation; mostly metamorphosed supracrustal rocks.
  5. Metapelitic, semipelitic, pyroxenite, pyroxenite, hornblendite; ultramafic sheets, lenses and boudins.
- **pPb4** — **4.** Marble, calcisilicate with diopside, serpentine, olivine, wollastonite; coarse-grained, white weathering; pure and impure metasedimentary magnesian carbonate.
- **pPb3** — **3.** Orthopyroxene garnet gneiss; orthopyroxene magnetite gneiss; silicate facies meta-iron-formation.
- **pPb2** — **2.** Pyrrhotite pyrite paragneiss; sulphidic two pyroxene gneiss, sulphidic pyrrhotite; metamorphosed sulphidic supracrustal rocks.
- **pPb1** — **1.** Pelitic gneiss with garnet, biotite, sillimanite, cordierite, graphite; lesser marble, calcisilicate, quartzite; orthopyroxene granitoid gneiss; metamorphosed supracrustal rocks.

ARCHEAN AND PALEOPROTEROZOIC: 3000 to 1920 Ma

THELON OROGEN, Slave craton cover

- **ApPbc** — **BOOTHIA COMPLEX:** granite, granitoid gneiss; lesser paragneiss, metapelite, local marble, calcisilicate, pyrrhotite; syn-orogenic plutonic complex interlayered with supracrustal rocks.
  5. Orthopyroxene-bearing garnet gneiss, commonly retrograded and locally migmatitic; minor metabasite and garnetiferous metasedimentary gneiss.
  4. Granite; red coloured, massive textured.
  3. Granitoid gneiss; amphibolite facies rocks retrograded from granulite grade.
  2. Orthopyroxene granitoid gneiss, paragneiss.
  1. Orthopyroxene granitoid gneiss; sheets of granite, pegmatite; lesser pelitic gneiss, marble, calcisilicate, pyrrhotite.

Contacts

- Geological boundary: defined or approximate

Faults

- Fault: approximate
- Fault: approximate, showing downthrown side
- Fault: assumed

Strike-slip faults

- Dextral strike-slip: approximate
- Dextral strike-slip: assumed
- Sinistral strike-slip: assumed

Thrust faults

- Thrust fault: approximate, teeth indicate upthrust side
- Thrust fault: assumed, teeth indicate upthrust side

REFERENCES

Blackadar, R.G., Christie, R.L., and Taylor, F.C., 1967. Geology, Boothia Peninsula and King William Island, District of Franklin; Geological Survey of Canada, Preliminary Map 2-1967, scale 1:506 880.

de Freitas, T.A., Harrison, J.C., and Mayr, U., 1997. Sequence stratigraphic correlation chart of the lower Paleozoic Franklinian succession, Canadian Arctic Islands and parts of north Greenland; Geological Survey of Canada, Open File 3410.

Dewing, K. and Nowlan, G.S., 2004. Correlation chart of Cambrian and Ordovician stratigraphy, Arctic Islands, Nunavut, Canada. Geological Survey of Canada, Open File 1837, 1p.

Frisch, T., 2011. Geology, Precambrian geology of northern Boothia Peninsula and Somerset Island, Nunavut; Geological Survey of Canada, Open File 6051, scale 1:250 000.

Harrison, J.C., Mayr, U., Mifflett, D.H., Sweet, A.R., McIntyre, D.J., Eberle, J.J., Harington, C.R., Chalmers, J.A., Dam, G. and Nøhr-Hansen, H., 1999. Correlation of Cenozoic sequences of the Canadian Arctic region and Greenland; implications for the tectonic history of northern North America; Bulletin of Canadian Petroleum Geology vol. 47, no. 3, 1999, pages 223-254.p.

Mayr, U., Frisch, T., de Freitas, T., Okulitch, A.V., and Kerr, J.W., 2003. Geology, eastern Prince of Wales Island and adjacent smaller islands, Nunavut; Geological Survey of Canada, Map 2016A, scale 1:125 000.

Mayr, U., Brent, T.A., de Freitas, T., Frisch, T., Nowlan, G.S., and Okulitch, A.V., 2004. Geology of eastern Prince of Wales Island and adjacent smaller islands, Nunavut; Geological Survey of Canada, Bulletin no. 574, 88 pages, 1 CD-ROM.

Stewart, W.D., 1987. Late Proterozoic To Early Tertiary Stratigraphy of Somerset Island and northern Boothia Peninsula, District of Franklin, N.W.T.; Geological Survey of CanadaGeological Survey of Canada, Paper 83-26, 1987; 78 pages.

Stewart, W.D. and Kerr, J.W., 1984. Geology, Somerset Island South, District of Franklin; Geological Survey of Canada, Map 1595A, scale 1:250 000.

Stewart, W.D. and Kerr, J.W., 1984. Geology, Somerset Island North, District of Franklin; Geological Survey of Canada, Map 1595A, scale 1:250 000.

Stewart, W.D. and Kerr, J.W., 1984. Geology, Boothia Peninsula, North, District of Franklin ; Geological Survey of Canada, Map 1597A, scale 1:250 000.

Stewart, W.D. and Kerr, J.W., 1984. Geology, Central Boothia Peninsula, District of Franklin; Geological Survey of Canada, Map 1598A, scale 1:250 000.

Thorsteinsson, R. (ed.) and Uyeno, T.T. (ed.)1980. Stratigraphy and conodonts of Upper Silurian and Lower Devonian rocks in the environs of the Boothia Uplift, Canadian Arctic Archipelago; Geological Survey of Canada, Bulletin 292, 75 pages.

Recommended citation
Harrison, J.C., Hillary, E.M., Ford, A., Stewart, W.D., Frisch, T., Mayr, U., and Thorsteinsson, R., 2013. Geology, Tectonic assemblage map of Peel Sound, Prince of Wales and Somerset islands, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 76 (preliminary), scale 1:500 000. doi:10.4095/292819

Preliminary Preliminary Preliminary

ess.nrcan.gc.ca

Canadian Geoscience Maps

Canada

Authors: J.C. Harrison, E.M. Hillary, A. Ford, W.D. Stewart, T. Frisch, U. Mayr, and R. Thorsteinsson

Geological compilation by J.C. Harrison

Source map geology (senior authors) by W.D. Stewart, T. Frisch, U. Mayr and R. Thorsteinsson

GIS development by T. Lynds

Spatial data capture by Gismo Solutions Ltd. (Edmonton) and A. Ford

Cartography by M.J. Baldock

Critical review by T. Hadlari

Initiative of the Geological Survey of Canada, conducted under the auspices of the Tri-Territorial Project as part of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) program.

CANADIAN GEOSCIENCE MAP 76

GEOLOGY

TECTONIC ASSEMBLAGE MAP OF PEEL SOUND

Prince of Wales and Somerset islands, Nunavut

1:500 000

10 0 10 20 30 40 km

Map projection Lambert Conformal Conic, standard parallels 71°30'N and 73°30'N. North American Datum 1983

Base map at the scale of 1:250 000 from Natural Resources Canada, with modifications.

Proximity to the North Magnetic Pole causes the magnetic compass to be erratic in this area.
Mean magnetic declination 2013, 19°02'W, decreasing 12.1'E annually. Readings vary from 4°09'W in the SW corner to 33°36'W in the NE corner of the map.

The Geological Survey of Canada welcomes corrections or additional information from users.

The data may include additional observations not portrayed on this map. See documentation accompanying the data.

This publication is available for free download through GEOSCAN (http://geoscan.nrcan.gc.ca/).

This map is not to be used for navigational purposes

Preliminary publications in this series have not been scientifically edited.

Preliminary Preliminary Preliminary

CANADIAN GEOSCIENCE MAP 76

GEOLOGY

TECTONIC ASSEMBLAGE MAP OF PEEL SOUND

Prince of Wales and Somerset islands, Nunavut