The innermost shelf off the Avalon Peninsula was completely unexplored in terms of geophysical surveys and bathymetric surveys until a summer 2009 GSC-A expedition (2009044). A series of surficial geology-related maps (panel at right) derives from the

opographic shaded-relief renderings generated (pre-expedition)

from Canadian Hydrographic Service spot depths significantly

enhanced topographic features and indicated a mix of bedrock

lineations, variable sediment cover, and small basins with some

sediment cover. While much of Newfoundland's inner shelf has been

examined with respect to glaciation, sediment deposits and glacial

chronology (Shaw 2002), the Avalon area remained largely

This is a unique terrain, contrasting with the relatively flat-lying, till-

blanketed Avalon Channel, to the east. The more rugged topography

of the late Proterozoic tectonized metasediments extending several

kilometres from land creates fjord-like bays, fault-governed ridges

and isolated basins. This contrasts with less indurated and more

flatlying Lower Paleozoic age metasediments which floor Avalon

The area was transgressed by the rising post-glacial sea-level and

would have bounded a proto-Labrador Current offshoot through the

adjacent Avalon Channel. A paucity of Quaternary deposits on land

that might afford dating potential has restricted understanding of the

late glacial history and it was suspected that the basins and harbours

might provide a datable sequence and a record of lower sea-level.

One goal was to explore for datable sediments in, for example,

isolation basins, marking the retreat of the ice sheet and the

terrestrial to marine change. Glacier flow directions, ice margins and

address. The potential for a gas pipeline landing, originating from

the Jeanne d'Arc Basin, was also a driver; this seascape had never

Multibeam bathymetric data across the Grand Banks is sparse, so

simply establishing geo-feature geometries (eg. bedrock and

sedimentary bedforms) requires considerable survey and

Hydrographic Service were accessed and gridded at appropriate bin

size for point spacings between 50 and 500 m in order to derive a

digital elevation model (DEM), a contour map, and a colour-coded

shaded-relief image of the seabed. This involved manual digitization

of one area, to fill a data gap, involving manual digitization of over 40

000 points. The DEM (panels at left) provided the control for survey

targets and for extrapolation and interpolation of map unit

low-stand apron

Prograding strata in lowstand apron

generally coincides with till or bedrock surface but

limited penetration precludes their differentiation

Distal glacimarine surface

Base of glacimarine;

Bedrock surface

Strata in bedrock

JD254/22:00


Proximal glacimarine surface

been characterized in terms of geology and morphology.


timing and the origin of Avalon Channel were also element to

channel and provide the platform for a large part of Grand Bank.

new survey and related findings are presented here.



Azig-zag pattern geophysical survey (expedition 2009044) was conducted to characterize the geology of the innermost shelf off the Avalon Peninsula (upper panel) and identify the past and present seabed processes. Detail of the sample stations visited in the Cape Broyle Harbour area of the Avalon south shore (lower panel). Rectangles correspond to



19) located just beyond the seismically homogeneous debrite lense. Core 19 penetration was sufficient to reach the base of the

glacimarine section but recovery was limited. At the base of the core, gravelly intervals at this base of slope setting represent either

intensification of IRD from the Labrador Current, the local affects of transgression across the bedrock ridge at the top of slope, or

alternatively, very small MTD activity. Core 16 had full recovery, shows evidence of the seismically-identified MTD debrite (see Cores

panel, to upper right) and enabled C-14 dating of the event.

## Geological conditions off the Avalon Peninsula, offshore easternmost Newfoundland: Bedrock and Glacial Features, Deglaciation Pattern and Chronology, Mass Failure and Attributes and Constraints to Engineering


Edward (Ned) L. King, Geological Survey of Canada-Atlantic, 1 Challenger Drive, P.O. Box 1005, Dartmouth, NS. eking@nrcan.gc.ca


shallow geology map units

shown in the Geologic

composite derived from high and mid-resolution seismic

(course alteration)





section and a disorganized shelly-pebbly unit whose derivation is unclear. At this time, nearby transgression was underway, supplying erosion

strata

JD255/00:00

his interpretation from a combination of sleevegun and deep-towed boomer outlines the general stratigraphy and morphologic features. The lower scale allows correlation with position on the ship's track in the far left panel.

products, the Avalon branch of the proto-Labrador Current was likely in operation, and small mass failure may still be occurring.

### The innermost shelf of the Avalon Peninsula is characterized in terms of Seismic reflection profiles together with two sediment cores helf bedrock geology, Quaternary sediment sequence, thickness and characterize the de-glacial and post-glacial sequence. Post-till deposition distribution, surficial texture, and glacially-related features. The latter include was followed by a stratigraphic succession of three mappable units, similar overdeepened and partially filled basins, drumlins and moraines, in depositional style but with a trend from poorly stratified to well stratified. deglaciation pattern and timing. The recent survey also allowed recognition both seismically and in the cores. The lowermost is barren of shells and of a paleo-mass transport event, lower sea-level indicators, sediment spurious investigation revealed few foraminifera, some reworked and some mobility under currents, iceberg scour distribution, including a recent event, typical glacimarine species. It is interpreted as meltwater-influenced, with and a spatial representation of quantified seabed roughness. Higher elevation and relief on the innermost shelf compared with the Avalon basin flank just outside Cape Broyle Harbour. Timing of retreat to the Channel reflects the bedrock change from Lower Paleozoic low-grade headlands is roughly constrained by the cores, with extrapolated metasediments below till in the Channel and harder, more deformed Late radiocarbon ages of about 15 ka (calibrated) at the till-glacimarine mud

Bedrock and Surficial Geology

Glaciations carved small valleys normal to the dominant N-S tectonic fabric Post-glacial of the bedrock which locally but rarely breached N-S offshore ridges and created locally overdeepened basins. Thick and thin till provinces were left, Recent seabed iceberg scouring is evident in sidescan images and seabed the thicker in shallow basins on the Avalon Channel floor (outside this area, photography. The frequency and magnitude of impact of relatively small the till is generally only 1 to 5 m thick) and the thinner, more patchy on the icebergs in shallow water and the location, magnitude and conditions of sand bedrock ridges areas. The offshore orientation of the drumlins and moraines transport are both largely unknown and relevant to pipeline or shows that the last vestiges of glaciation emanated from the Avalon ice communications cables. Initial observations suggest both processes are dome, traversing normal to the Avalon Channel. It was vigorous enough to locally active. generate drumlin fields but on retreat the ice sheet became thin enough to be sensitive to local topography; it assumed a lobate regime locally, reflected in **Engineering** trends of transverse (Rogen-like) moraines but while still streaming in the

The innermost shelf of the Avalon Peninsula, like the coastline, has a high

relief, where morphology is dominated by Proterozoic bedrock with strong

structural elements. Yet glacial erosion has locally cut offshore extensions

of some harbours, often with overdeepening. The offshore contact with

A series of surficial geology maps complement this poster (separate

product) and include 3-dimensional information, such as sediment thickness

the younger (lower Paleozoic) and more erodible rock sequence

coincides with a till-dominated terrain.

sedimentation rates (>4 metres per kyr).

Sea-level Low-stand Deposit

and Quaternary stratigraphic succession distribution.

A small (1 million cubic m) debrite is dated about 14.8 ka; much smaller event night have occurred as late as 12 ka. Thick low-stand deposits were likely fed by coastal glaciers at stands about 45 and 35 m present water depth. These help constrain the regional

Previously unknown glacially-cut valleys are rare but afford potential pipeline

routes where issues with rugged terrain, difficult trenching materials, shallow

some IRD emanating from the glacial margin while it was positioned within

the catchment of the basin. At some point, the margin likely lay at the SW


mass failure occurred, at about 14.8 ka (cal.). This was followed by retreat


from the small basin to the harbours where it deposited glacifluvial or ice

contact deltas, probably under the influence of a rising relative sea-level.

The survey covered areas too deep to identify isolation basins (lakes drowned with post-glacial sea-level rise) yet small examples may be present. Sand mobility under current influence periodically activates thin bedforms.


Drumlins and moraines indicate a relatively thin and lobate ice configuration A modern and relict iceberg scour population is recognized with fresh scours The overdeepened basins are the site of glacimarine sediments that A glacial imprint on the Avalon Peninsula east coast presents elements recorded deglaciation initiating at about 15 to 14.8 ka with high introducing constraints and mitigating opportunities for avoidance of difficult terrain for routing, burial and landfall of any potential gas pipeline.





Proterozoic metasediments inside the ca. 150 metre isobath, about 5 km off contact. Soon after, the basin was inundated by the ocean at least one small

| Expedition-<br>Core No. | Sample<br>Interval<br>cm | Species or<br>Genus ID | GeoSetting   | Conven-<br>tional<br>C-14 (BP) | Calib.*<br>Median,<br>BP | Calibrated*    |                    | Lab. &              |
|-------------------------|--------------------------|------------------------|--|--------------------------------|--------------------------|----------------|--------------------|---------------------|
|                         |                          |                        |  |                                |                          | Range,<br>BP   | Range,<br>Calendar | No.                 |
| 2009044-0016            | 95 - 97                  | Yoldiella sp.          | mid-section of the uppermost late glacimarine unit (Gmws)  | 13920 +/- 70                   | 13954                    | 14393 to 13626 | 16343 to 15576     | Beta-<br>276646     |
| 2009044-0016            | 163 - 165                | Yoldia<br>limatula     | top of the uppermost late glacimarine map<br>unit (Gmws); matches the top (immediately<br>above) of the seismic-defined debris flow                        | 13900 +/- 70                   | 13911                    | 14526 to 13252 | 16476 to 15202     | Beta-<br>276655     |
| 2009044-0019            | 92 - 94                  | Geukensia<br>demissa   | small basin flank; seismic shows several m<br>well stratified fill (map unit Gmws) on SW<br>basin flank; this is highest stratigraphic level<br>with shell | 8120 +/- 50                    | 6239                     | 6341 to 6168   | 8291 to 8118       | Beta-<br>276647     |
| 2009044-0019            | 164 - 166                | unidentifiable         | core intersects stratified shell-rich debris near<br>base; transgression related storm, tsunami or<br>debris flow deposit; very shell-fragment rich        | 9280 +/- 50                    | 7593                     | 7666 to 7514   | 9616 to 9464       | Beta-<br>276657     |
| 2009044-0019            | 248 - 250                | unidentifiable         | core likely does not reach debris flow horizon;<br>coastal shells transported with transgression<br>related storm, tsunami or late debris flow             | 10700+/- 25                    | 9330                     | 9383 to 9258   | 11333 to 11208     | UC Irvine-<br>87625 |



# **Engineering-Related Characterization**

This study was partly aimed at investigating the innermost shelf in terms of bedrock, terrain Meso-scale seabed roughness from Huntec DTS sounder profiles was calculated through four and Quaternary geology that might be encountered should a gas pipeline be routed directly from the hydrocarbon production areas on Grand Bank. A seabed roughness characterization displays columns proportional to elevation deviations from four smoothed classes (technique to right). Relatively soft sediment-filled glacially excavated channels afford potential pipeline routes leading to shoreline cliffs, should it be advantageous to avoid adjacent high relief hard

Seabed photographs in a

Cape Broyle Harbour megaripple field just

south of Brigus Head (stations 17, camera and

20-21, Van veen grabs).

The megaripples are very

wavelength, and manifest

as a veneer of sand on a

gravel lag which supports

considerable biota. The

grab samples yielded a well sorted fine dark grey

of assorted lithologies. A

usting of flock on the fine

to medium sand suggests


current activity is not only

tidally-generated and that

storm-related currents

may be influential.

The two panels below are enlargements of maps depicting thickness and distribution of the



## **Calculation of Seabed Roughness**

Seabed photographs (Stn. 18) across a bedrock, till, mud and iceberg scoured terrain immediately seaward of

looking ice scours occurs on the till-covered hummocks of the area; those in the basins are more sediment-filled.


Cape Broyle Harbour (location, panel far left). The background is a low-resolution bathymetric DEM with contours


cobbles. A dusting of flock covers the seabed; photo 19 is an exception where iceberg scouring has disturbed

and superimposed sidescan imagery showing a variety of features. The mud is very thin, just covering gravel and


coarse gravel, though fishing activity may also be responsible. An apparently disproportionate number of fresh-

smoothing classes. A guided autopick of the seabed (ie. bathymetric readings) was first smoothed to remove wave-action created heave using a GSCA-developed heave filter program (DeJitter-P. Pledge). The other colours represent progressively greater filtering (running averages) to establish different seabed trends. The deviations from these trends are then calculated and plotted as columns superimposed on the 3-D topography rendering. Meso-scale roughness represents the irregularity attributed to seabed features such as bedrock, iceberg scours, pockmarks, buried channels, drumlins and smooth sandy or muddy seabeds.












The smaller, superimposed ridges on the boomer profile (C) are marked by blue arrows and correspond to the ribbed moraines on the sidescan. They are up to 5 m relief and typically ~50 to 75 m across with spacings of 75 to 150 m. The largest ridge is 200 m across and asymmetric in profile. A larger field of the features is situated in a sub-basin of the Avalon Channel, Box E, each interpreted as ribbed (transverse) moraines (similar to Rogen or deGeer potential challenge to pipeline routing. moraines). Box G shows their distribution.



aligned with Huntec profiler section

glacimarine sediment, marking the top of till. Box F shows selected seabed are of similar relief to "megafurrows" (iceberg scours), an example of which is located NE of Cape Race (Fader 1989). It is unlikely photographs from a transect across several ridges as identified from that the modern iceberg population is capable of "megafurrows"; the "sill" to the north within Avalon Channel would have a filtering sidescan. They are cobble and gravel-topped with a thin sand cover on the till effect on the largest ones such that it is likely that this megafurrow derived from one of many relict icebergs calving directly from the and possibly limited glacimarine ponded between the ridges. They are nearby (paleo) coast. Still, both features, with cobble-rich crests and clast-rich, high-strength, cohesive interstitial clay, present







Relief in the DEM is too sparse to show drumlin orientation but the geophysical data combined with the DEM spot elevations allow determination of their

Both the drumlins and subsequently-formed moraines indicate that at least the latest glacial ice imprint on the seabed flowed from the Avalon Cap and not along the Avalon Channel. This glacier must have been quite rigerous as drumlins are generally associated with streaming ice. The swing in moraine orientation (partly schematic yellow lines, box A) generally follows the contour trend of a bathymetrically-defined lobate body. This indicates that the glacier was thin enough for this subtle sub-glacial form to affect local flow direction and a lobate (slightly radially flowing) ice tongue developed during deglaciation. As no stratigraphic differentiation is visible in the thick till it is not possible to deduce any regional evolution in flow direction. The Avalon Channel, a large geomorphic feature, may have formed entirely through flow-transverse erosion (overdeepening). The pre-Cambrian rock clasts at it's sole are harder han the offshore bedrock, affording a mechanism for preferential erosion at and down-ice from the bedrock contact. approximate boundaries but only along the survey tracks. The white hatch

The surficial map compilation and seismic section picking was by COOP student Nader Mostaghimi and the author. Core analysis was by Marla Reid my Pellerin, Albert Rand and Midori Tellus-Langdon under the direction of Jenna Higgins and Kate Jarrett, GSC-A core facility. Radiometric carbon dates were by the Beta Analytic AMS Radiocarbon Dating Lab, FL, with the exception of one analysis from the Keck Carbon Cycle AMS Facility, CA (core 19; 249 cm depth). Seismic compilation benefitted from the software tools and ESRI ArcMap interactivity developed by Bob Courtney and Paul Fraser, GSC-A. The crew and Captain Naugle of GSC-A expedition 2009044 aboard CCGS Hudson and the invaluable effort of numerous GSC-A scientific and technical staff made data collection possible. This project was supported by the GSC GECOD project (Geoscience for East Coast Offshore

Development) under Michael Li's direction, and PERD funding. Gordon

Cameron provided valuable critical review and scientific discussion.

outlines the field. Moraines, as traced from sidescan images show a dominant

**Drumlin and Moraine Configuration** 

-S orientation, normal to the drumlins.

# Fader, G.B.J., 1989. Submersible observations of iceberg furrows and sand King, E.L., 2013. Geological conditions off the Avalon Peninsula,

ridges, Grand Banks of Newfoundland in: Submersible observations of the east coast of Canada, (ed.) D.J.W.Piper; Geological Survey of Canada, Paper 88-20, 27-39.

Shaw, J., 2003. Submarine moraines in Newfoundland coastal waters: implications for the deglaciation of Newfoundland and adjacent areas. Quaternary International 99-100, 115-134.

Shaw, J. & Forbes, D.,1995. The postglacial relative sea-level lowstand in Newfoundland. Canadian Journal of Earth Science, 32, 1308-1330.

This publication is available for free download through GEOSCAN (http://geoscan.ess.nrcan.gc.ca/).

Attributes and Constraints to Engineering: Geological Survey of Canada, Open File 7360 (revised), Poster. doi:10.4095/292863 DOSSIER PUBLIC have they

Features, Deglaciation Pattern and Chronology, Mass Failure and

offshore easternmost Newfoundland: Bedrock and Glacial

Recommended citation

OPEN FILE Publications in the **7360** (revised)

2009044: Station 22 Camera Stills

superimposed on Klein sidescan image Camera Transit; Ship's antenna; start and end JD/UTC time

Photo locations;

