

DESCRIPTIVE NOTES

The map area (NTS 87-114) lies within the Minto Inlier, a ~300 km long by 100–150 km wide belt of gently folded sedimentary and igneous rocks of early Neoproterozoic age (Tonian-early Cryogenian) age. The Neoproterozoic sedimentary strata belong to the Shaler Supergroup, an approximately 4 km thick succession of shallow marine carbonate rocks and evaporite rocks with interbedded terrigenous rocks that were mainly deposited in a shallow intracratonic septic sea, referred to as the Amunden Basin (Rainbird et al., 1994; Rainbird et al., 1996a; Thorsen and Tozer, 1992; Young, 1991). The basin is considered to have formed within the supercontinent Rodinia and exposures of similar rocks in what are now the Mackenzie Basins of the northern Cordillera suggest that it extended for more than 1000 km to the southwest (Long et al., 2006; Rainbird et al., 1996b). The sedimentary succession is intercalated with mafic sills of the ca. 720 Ma Franklin igneous province (Hearn et al., 1992). The sills of variable thickness up to more than 100 m thick in the south. In many cases, individual sills extend for 20 km or more along-strike with little significant change in thickness. Sills constitute anywhere from 10 to 50 per cent of the stratigraphic section. Sills of similar type and age also occur in the Coppermine terrones, Brook Inlet and Duke of York Inlet to the south (Rainbird et al., 1996b; Steinhilber et al., 2004) and are geochemically similar intrusions and volcanic rocks associated with the Franklin event extend from Greenland to the western Yukon (Derynck et al., 2006; Hearn et al., 1992; Macdonald et al., 2010). The Shaler Supergroup member is capped by a succession of flood basalts flows and interflow sedimentary rocks (Nauyasik Fm.) more than 1 km thick, which are the extensive equivalent of the northern Cordillera (Rainbird et al., 1996b). Rare north-west-trending dykes are interpreted to have intruded along syn-magmatic normal faults, to feed sills and possibly the flood basalts (Bedard et al., 2012). The stratigraphic sequence is identified in the lower units of the lower section (East side of the Shaler Supergroup). The oldest sills and corresponding basal lavas are enriched in incompatible trace elements and may have silvite-enriched bases. These mafic sills correspond to the major sheet flow units of the lava succession (East side of the Shaler Supergroup). The sills are exposed only at the north-west end of Minto Inlier, near Hadley Bay, where they unconformably overlie Paleoproterozoic sedimentary rocks, which in turn unconformably overlie Archean granitic rocks (Campbell, 1981; Rainbird et al., 1994). The irregular edge of Minto Inlier is defined by an essential unconformity that separates the Neoproterozoic rocks from Lower Cambrian sandstones and siltstones that pass upward into a thick succession of mainly siliceous carbonate rocks, ranging in age from Cambrian to Devonian (Thorsen and Tozer, 1992). Structurally, the Minto Inlier is relatively simple, composed of the open, north-trending Haiman Island syncline and a smaller Walker Bay anticline to the northwest. Beds typically dip less than 10° and there is generally no penetrative cleavage or other apparent outcrop-scale fabric. The origin of the folding is unknown but it occurred after deposition of the early Neoproterozoic rocks and before uplift, erosion and deposition of overlying lower Cambrian siliceous rocks, which are not folded. All rocks are dissected by east-northeast to east-trending faults that form a horst and graben system with up to 200 m of normal displacement on individual faults. The zones of faulting is approximately 100 km wide and stretches from the head of Minto Inlet in the east and is spectacularly imaged as prominent lineaments on recently published aeromagnetic maps (e.g. Kiss and Orneschko 2010).

NTS 87-114 is underlain by stratigraphic units from the middle to upper Kilian Formation, Kuujuaia Formation and Nauyasik Formation of the Shaler Supergroup. Together with diabase sills, the strata comprise the gently south-dipping northern limb of the Haiman Island Syncline, whose axis lies along the southern edge of the map sheet. Exposures of the Kilian Formation (clastic-carbonate) and tan carbonate members are limited in the northern half of the map sheet area due to the massive nature of the strata and the dominance of thick diabase sills that form a prominent southeast dip slope down to the Kuujuaia River. The upper Kilian Formation, Kuujuaia Formation and basal Nauyasik Formation, are best exposed along a prominent cuestas that faces north-northeast along the south side of the Kuujuaia River. The Kilian generally is quite resistant but a good exposure of the tan carbonate member (nPK3) and upper evaporite member (nPK4) is located near UTM 566071E, 786497N. In most places within the map area, the basal contact of the Kuujuaia with the upper evaporite member (nPK4) is very distinct, but at one locality (UTM 565163E, 7860759N), it is a well developed erosional unconformity overlain by a thin, discontinuous, chert-oxidized conglomerate. The conglomerate, in turn, is overlain by a green, parallel-stratified silty-sandstone, which passes upward into a dark green siltstone. This is interbedded with a cleaner quartzite sandstone, more typical of overlying crossbedded quartzite of the Kuujuaia Formation. The underlying Kilian Formation strata are also well exposed (chert-oxidized siltstone and sandstone), suggesting that deposition of the overlying strata was accompanied by introduction of reducing hydrothermal fluids. This important contact zone that volcanically normally associated with the overlying Nauyasik Formation, was already active at the beginning of Kuujuaia time. One of the thickest and best preserved sections of the Kuujuaia Formation is exposed near the eastern end of the cuestas at UTM 57140E, 786389N (see section W-22 of Rainbird, 1992 and section 4 of Jefferson, 1995). Other good exposures of the Kuujuaia Formation are located at UTM 56430E, 789989N (section 86-10) and UTM 564146E, 789497N (section 86-14). The Kuujuaia and Nauyasik formations are also well exposed along the flanks of two, north-flowing tributaries of the Kuujuaia River. The contact between the two formations generally is planar and conformable but there are numerous places where soft-sediment deformation features are preserved in the underlying sandstone indicating that the sand was un lithified (see Rainbird, 1995). As well, irregular fragments of basalt have been observed within the sandstone. The most complete exposures of the Nauyasik Formation are located at approximately UTM 56484E, 789502N and UTM 56481E, 789507N and are described in detail by Williamson et al. (2013). Members nPK1 and nPK2 are composed of basaltic flows and associated interflow deposits that are regionally distributed, but the intervening volcanoclastic nPK2 member is discontinuous and of variable thickness. It then marks the central part of the map area lowest in the northeast and is absent in the western half of the map area. At least 5 diabase sills occur within the map area and generally are spaced at regular intervals within the host sedimentary rocks. The sills are of the type 2 (diabase) described in the legend. Several steep, north-west-trending, mostly west-side down, normal faults are spaced evenly across the map area and are evident as prominent linear topographic lows. At approximately UTM 56451E, 786470N, the lineament dykes, suggesting that emplacement was coincident with faulting, a relationship that has been documented elsewhere in the region (Bedard et al., 2012). East-west-trending faulting, common in map areas to the north, is largely absent from NTS 87-114.



Abstract
NTS 87-114 is underlain by the middle to upper Kilian, Kuujuaia and Nauyasik formations of the Neoproterozoic Shaler Supergroup. Together with diabase sills (type 2), spaced at regular intervals within the host sedimentary rocks, the strata comprise the gently south-dipping northern limb of the Haiman Island Syncline. The northern half of the map area is dominated by a thick sill, which forms a prominent southeast dip slope down to the Kuujuaia River. Sedimentary strata are best exposed along a prominent cuestas that faces north-northeast along the south side of the Kuujuaia River. The upper two members of the Kilian Formation are especially exposed along the base of the cuestas crossbedded quartzite of the Kuujuaia Formation. The overlying Kuujuaia Formation is prominent along the west, north-flowing tributaries of the Kuujuaia River. Several steep, north-west-trending, west-side down normal faults are spaced evenly across the map area.

Résumé
Le feuillet NTS 87-114 expose des roches Néoproterozoïques des Formations du Kilian Moyer à Supérieur, Kuujuaia et Nauyasik. Les roches sédimentaires appartiennent au Supergroupe de Shaler, et sont injectées par 5 flows-carbonés, diabasiques de Feilermment Franklin. Les strates constituent le flanc nord du Synclinal Haiman Island, et sont dominées vers le sud. La partie nord du feuillet est dominée par une énorme masse de basalte, qui forme un banc ou va jusqu'à la rivière Kuujuaia. Les roches sédimentaires sont bien exposées le long d'une falaise qui s'élève au sud de cette rivière et assure le long de ses tributaires du sud. Les deux membres supérieurs du Kilian sont exposés séparément au pied de la falaise. Les arénites quartzitiques à lits entrecroisés de la Formation de Kuujuaia sont en discordance sur les strates du Kilian. Le sommet de la montagne est formé par les lavas et volcanoclastiques de la Formation de Nauyasik. Il y a plusieurs failles normales orientées nord-ouest, et une pendage vers l'ouest.

Cover illustration
Looking east at flood basalt-capped cuestas with Kuujuaia River and upper Kilian Formation and diabase sill in foreground, Victoria Island, Northwest Territories, 2011-07-13. Photograph by R. Rainbird, 2013-204.

Catalogue No. M183-1104-2012E-PDF
ISBN 978-1-106-21440-5
doi: 10.4095/29344

© Her Majesty the Queen in Right of Canada 2013



CENOZOIC
Q Quaternary sediments.

NEOPROTEROZOIC
nPK1 Franklin intrusions: Typically massive, laterally extensive, diabasic sills with columnar jointing (~50 m thick, rarely up to 100 m). Some sills are composite with internal intrusive contacts. Two types: 1) In older, more mafic sills, the columnar jointing is more regular and the sills are more mafic. The sills are capped by a thin (1–2 m) hydrothermal pyroclastic tuffaceous. The dioritic cumulate is commonly covered with bright orange lichen, weathers chocolate brown, and shows a characteristic layer-parallel ribbed weathering. Upper 1/3 to 1/2 of sills composed of massive olivine and pyroxene gabbro, a magnetite gabbro with common pitted weathering (magnetite ochreous) and a granophyric horizon containing abundant ocelli of granophyre and coarse, tabular oligoclase crystals. 2) Younger (based on cross cutting relationships), more evolved, diabasic sills showing enrichment in magnetite, ilmenite, quartz and alkali feldspar towards their cores, but are rarely layered. Some sills are porphyritic and contain 10–15% plagioclase-oligoclase-epidote-epidote phenocrysts and granophyre up to 5 mm. Less common, 1–40 m wide dykes, irregular to very linear (generally oriented NNW). Commonly associated with fault breccias or drag folds in host metasediments. Dykes commonly connect to sills, some associated with calc-silicate contact metamorphic rocks (reddish garnet overlain by bright green vesicular), black Fe-oxide skarns, and minor sulphides.

nPK2 Nauyasik Formation (nPN1–nPN3)
Sheet-flow member: Bluish-green to orange-weathering, laterally extensive, subaerial basalt flows. Individual flows 15 to 50 m thick. Flow structure varies from columnar-dike-like to a massive base with typically vesicular flow tops. Rare interflow scoria, spatter, fumarolic concretions, volcanic necks and pillis to disseminated native copper. Maximum thickness of 200 m, limited by erosional preservation.

nPK3 Lower receiver member: Massive to parallel-stratified volcanic-pebble conglomerate overlain by thin, parallel to cross-bedded, quartz-rich volcanoclastic sandstone. Framework composed of volcanic, carbonate and sandstone clasts, matrix rich in quartz sand grains. Conglomerate sub-unit varies in thickness from 40 to 100 m, sandstone sub-unit from 5 to 10 m.

nPK4 Lower member: Dark green to grey weathering, dominantly subaerial flows, varying from fine massive basalt to coarse sub-ophitic basalt. Filled and types sand breccias are common at unit base indicating emplacement into shallow water. Thin (1 to 10 m) sheet flows with massive bases and vesicular flow top, or discontinuous lobate flows. Degree of vesicularity varies throughout. Thickness 40 to 70 m.

nPN1 Shaler Supergroup (nPK1–nPK4)
Kuujuaia Formation: Two principal lithologies: coarse quartzite composed of tabular coarse-grained siltstone and compound planar crossbedding and a less abundant fine-grained assemblage of interbedded fine sandstone, dolomite and mudstone forming lenses up to 20 m wide. Rare basaltic pebbles, ~120 m thick.

nPN2 Kilian Formation (nPK1–nPK4)
Upper Evaporite-Carbonate member: Base is dolomite and dolomite with 10–20% ripple crossbedded siltstone. Bedding parallel and crosscutting salt-pipe ventrites and desiccation cracks common. Changes up-section from creamy grey to pinkish grey, reflecting increase in hemalitic siltstone relative to carbonate. Nodular sulphate more common in middle part of member. Upper consists mainly of parallel-laminated red dolomite mudstone and waxy to ventricular bedded, buff to pink-weathering dolomite (no sulphate). Diagenetic redox horizons, desiccation cracks, halite pseudomorphs and trace structures are ubiquitous. Present only on the southwest domain of the Minto Inlier, ~80 m thick.

nPN3 Tan Carbonate member: Tan to green-grey, flaggy weathering dolomite and limestone. Gradation between parallel-laminated lute and flat to waxy and hummocky bedded siltstone. Lute-rich layers are generally plane parallel scoured grading up to lute-rich type. Transformational clast breccia commonly infilling swales and gutters. Black chert nodules throughout and sporadically at several horizons. One distinctive bedform, from the middle of the tan carbonate member, is laterally traceable from Ukhahkak along the Kuujuaia River valley to where it cuts across the Nauyasik plateau, ~60 m thick.

nPN4 Clastic-carbonate member: Variegated (red, green, grey and black) pin-stripe laminated mudstone and siltstone, particularly at its base. Desiccation cracks common in mudstone and waxy basing; ripple crossbedding in coarse siltstone-fine sandstone interlayers. Waxy-flaser bedded and small-scale crossbedded, ~4 m thick, buff-weathering, fine-grained quartzites near top. Wavy bedded dolomite and laterally linked stromatolite interbeds are common and increase up-section, ~120 m thick.

nPN5 Carbonate-evaporite member: Alternating, decimetre-scale subunits of evaporite carbonate-dominated lithologies, evaporite laminated red mudstone and dolomite mudstone with interbedded nodular arylthrite and laminated opaline and arylthrite, minor stromatolite dolomite. Carbonate lithologies: dolomite and minor limestone lute-rich rhythmic capped by arylthrite laterally linked stromatolites, forming repetitive metre-scale cycles. Meter-tooth structure common.

Contact, depositional, depositional-conformable or intrusive
Defined
Approximate
Inferred

Contact, depositional-unconformable
Defined
Approximate
Inferred

Fault, generic, steep dip
Defined
Inferred

Fault, normal, upright
Defined
Approximate
Inferred

Fault, reverse, upright
Defined
Approximate

Structural lineament
Defined
Approximate

Station location
Ground observation

Planar structure
Fault
Vertical
Inclined
Horizontal

Foliation
Igneous layering

Linear structure
Lineation
Fault strike, fault grooves, slickensides, or mineral growth fibres
Glacial station or groove