


Characterization of mineralizing fluids and fluid-structural relationships in the Phoenix uranium deposit, southeastern Athabasca Basin, northern Saskatchewan: preliminary core and petrographic studies

K. Wang¹, G. Chi¹, K.M. Bethune¹, Z. Li¹ and C.D. Card²



Introduction

The Phoenix deposit, a high-grade unconformity-related uranium deposit associated with a NE-trending moderately SE-dipping reverse fault (WS shear zone), is located 35 km southwest of the world-class McArthur River uranium deposit in the southeastern Athabasca Basin (Fig. 1). Previous geological and geochemical studies have revealed features similar to other unconformity-related uranium deposits in the region (e.g. Kerr, 2010; Power et al., 2012; Dann et al., 2014), but the composition, temperature and pressure of the fluids associated with mineralization have not yet been investigated with fluid inclusion techniques. Furthermore, like other well-studied unconformity-related uranium deposits, the structural controls on mineralization, particularly the hydrodynamic relationship between structures (and related stresses) and fluid pressure, are not well understood. The main objectives of this study are to try to determine the fluid composition, thermal conditions and fluid-structural relationships in the Phoenix deposit through a combination of field (drill-core) investigation and petrographic, microstructural and fluid inclusion studies.

Location




Fig. 1: The Phoenix deposit is geographically located in the southeast corner of the Athabasca Basin, 35 km southwest of the McArthur River mine and 25 km northeast of the Key Lake mill (modified from Card et al. 2007).

Field drill-core investigation




Fig. 2: Map locations of studied drill cores, WR-260, WR-440, WR-250, and WR-394 (modified from Arseneau and Revering, 2010)




Fig. 3: Schematic cross section of the Phoenix deposit, modified from Gamelin et al. (2010)

- Thirty-five samples were examined from the six drill cores (Fig. 2) to determine lithological characteristics and document different varieties of fracture/vein fillings and associated alteration; six of the samples are oriented (Fig. 4-B&D);
- The basement is dominated by granulite grade graphitic pelitic to psammopelitic rocks (Fig. 4-A);
- The pelitic rocks are invaded by dm to m-thick foliation parallel bodies of massive quartz (Fig. 4-B) and spatially associated (gradational) pegmatite (Fig. 4-C), in places with extensive silicification of host rocks;
- Graphite of probable hydrothermal origin is abundant in these zones (Fig. 4-D);
- Intense silicification was observed along the unconformity surface, and clasts of 'quartzite' (hydrothermal quartz?, Q-H) were observed in the basal conglomerate of the Athabasca Group (Manitou Falls Fm [Read Fm in Fig. 3]);
- A younger generation of fractures and microbreccia zones, characterized by vuggy quartz infillings, cut both the silicified basement rocks (Fig. 4-E) and overlying sandstone (Fig. 4-F); Locally these fractures are lined by graphite




Fig. 4-A: Garnetiferous pelite (WR-440)



Fig. 4-B: Oriented sample of hydrothermal quartz (WR-440)



Fig. 4-C: Pegmatite cut by graphite-filled fractures (WR-440)



Fig. 4-D: Hydrothermal graphite vein cutting granite (WR-440)



Fig. 4-E: Pelite with hydrothermal quartz cut by vuggy quartz vein (WR-440)




Fig. 4-F: Vuggy Quartz cutting sandstone (WR-250)

Petrographic study




Fig. 5-A: Hydrothermal quartz ('quartzite', Q-H, grain size up to 28mm)




Fig. 5-B: Numerous needle-like mineral inclusions (rutile or tourmaline?) in Q-H




Fig. 5-C: Garnet-sillimanite bearing pelitic gneiss




Fig. 5-D: Q-H cutting garnet-cordierite-sillimanite-bearing pelitic gneiss




Fig. 5-E: Needle-like mineral inclusions in a Q-H vein




Fig. 5-F: Sudoite and muscovite alteration




Fig. 5-G: Vuggy quartz (Q-V) cutting quartz conglomerate




Fig. 5-H: Q-V cutting quartz sandstone






Fig. 5-I: Q-V cutting Q-H



Fluid inclusion (FI) study





Figs. 6-A&B: CO₂-dominated inclusions are well developed in densely distributed, parallel microfractures in the hydrothermal quartz, which predates mineralization.



Figs. 6-C, D, E&F: abundant vapour-only aqueous inclusions found in the euhedral crystals of the secondary vuggy quartz veins and cements.



Figs. 6-C, D, E&F: abundant vapour-only aqueous inclusions found in the euhedral crystals of the secondary vuggy quartz veins and cements.



Summary and conclusions

- Collectively, the field and petrographic observations, coupled with preliminary fluid inclusion analysis, are consistent with a two-stage history:
 - extensive pre-Athabasca hydrothermal alteration with associated generation of hydrothermal quartz (Q-H) and pegmatite;
 - post-Athabasca brittle faulting/fracturing and infiltration of a second generation of quartz (Q-V);
- Petrographic studies of the quartz-rich rock (Q-H) indicates that the quartz grains are very coarse (up to 28mm) and contain abundant needle-like mineral inclusions (rutile or tourmaline?), suggesting the rock may be quartz generated by hydrothermal activity;
- CO₂-dominated inclusions are well developed in densely distributed, parallel microfractures in the hydrothermal quartz (Q-H) in the basement, suggesting relatively high fluid pressure;
- Abundant vapour-only aqueous inclusions were found in the euhedral crystals of the secondary cements and vuggy quartz veins (Q-V) near the unconformity and in the overlying Manitou Falls Formation. This, together with the low homogenization temperatures (101 to 134°C) of co-existing biphasic aqueous inclusions, suggests low fluid pressures and probable boiling;
- The brittle-style deformation associated with emplacement of this secondary quartz contrasts with the ductile deformation in the basement;
- At present it is uncertain whether or not the fluids represented by these inclusions were directly related to mineralization;
- Ongoing studies aim to characterize the nature and sequence of structures and related mineralizing fluids (fluid composition, temperature and pressure), as well as to evaluate the stress regime during mineralization, through the orientation of microfractures and fluid inclusion planes.

Acknowledgments

This project is supported by the TGI-4 (Targeted Geoscience Initiative Four) program of Natural Resources Canada. The study has benefitted from collaboration with the Saskatchewan Geological Survey. We also would like to thank Denison Mines Corporation for supporting the field logistics.

References

Arseneau, G. and Revering, C., 2010. Technical Report on the Phoenix Deposit (Zones A & B) - Wheeler River Project, Eastern Athabasca Basin, Northern Saskatchewan, Canada. NI 43-101 technical report prepared for Denison Mines Corp. by SRK Consulting.

Card, C., Pana, D., Portella, P., Thomas, D.J., Amesley, I.R., 2007. Basement rocks of the Athabasca Basin, Saskatchewan and Alberta, in EXTECH IV: Geology and Uranium Exploration Technology of the Proterozoic Athabasca Basin, Saskatchewan and Alberta, G.W. Jefferson and G. Delaney (eds.), Geol. Surv. Canada, Bulletin 588, p. 69-87.

Dann, J., Hattori, K., Potter, E.C., and Sorba, C., 2013. Discrimination of Elemental Assemblages in the Alteration Halo of the Phoenix Deposit, Saskatchewan, Through Applied GIS, Geological Survey of Canada, Open File 7463, doi:10.4095/293122.

Gamelin, C., Sorba, C., Kerr, W., 2010. The discovery of the Phoenix Deposit: A new high-grade Athabasca Basin unconformity-type uranium deposit, Saskatchewan, Canada. Saskatchewan Geol. Surv. Open House 2010 Abstract Volume, p. 17.

Kerr, W.C., 2010. The discovery of the Phoenix deposit: a new high-grade Athabasca Basin unconformity-type uranium deposit, Saskatchewan, Canada, In R.J. Goldfarb, E.E. Marsh and T. Monecke (eds.), The Challenge of Finding New Mineral Resources: Global Metallurgy, Innovative Exploration, and New Discoveries, Volume II: Zinc-Lead, Nickel-Copper-PGE, and Uranium, Society of Economic Geologists, Special Publication 15, p. 703-725.

Power, M.J., Hattori, K., Sorba, C., and Potter, E.C., 2012. Geochemical anomalies in the soil and uppermost siliciclastic units overlying the Phoenix uranium deposit, Athabasca Basin, Saskatchewan, Geological Survey of Canada, Open File 7257, 36 p. doi:10.4095/291981