



Natural Resources
Canada

Ressources naturelles
Canada

CANADIAN GEOSCIENCE MAP 160

GEOLOGY

NIPPERS HARBOUR AND PARTS OF HORSE ISLANDS, CAPE ST. JOHN, AND LITTLE BAY ISLAND

Newfoundland and Labrador

NTS 2-E/13 and parts of NTS 2-E/12, NTS 2-E/14, and
NTS 2-L/4



Map Information Document



Canadian Geoscience Maps

Canada

PUBLICATION

Map Number

Natural Resources Canada, Geological Survey of Canada
Canadian Geoscience Map 160

Title

Geology, Nippers Harbour and parts of Horse Islands, Cape St. John, and Little Bay Island, Newfoundland and Labrador, NTS 2-E/13 and parts of NTS 2-E/12, NTS 2-E/14, and NTS 2-L/4

Scale

1:50 000

Catalogue Information

Catalogue No. M183-1/160-2013E-PDF
ISBN 978-1-100-22550-0
doi:10.4095/295866

Copyright

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources Canada, 2015

Recommended Citation

Skulski, T., Castonguay, S., Moussallam, Y., McNicoll, V.J., van Staal, C.R., and Bédard, J.H., 2015. Geology, Nippers Harbour and parts of Horse Islands, Cape St. John, and Little Bay Island, Newfoundland and Labrador, NTS 2-E/13 and parts of NTS 2-E/12, NTS 2-E/14, and NTS 2-L/4; Geological Survey of Canada, Canadian Geoscience Map 160, scale 1:50 000. doi:10.4095/295866

ABSTRACT

Northeastern Baie Verte Peninsula (Newfoundland and Labrador, NTS 2-E/13, parts of NTS 12-I/4, 2-E/14, and 2-E/12) is underlain by the Neoproterozoic to Ordovician Ming's Bight Group, Cambrian ophiolite of the Betts Cove Complex, submarine Ordovician ophiolite cover of the Snooks Arm Group, and a continental overlap assemblage including Ordovician-Silurian Burlington plutonic suite, Stocking Harbour intrusive suite, and the Silurian Confusion Bay plutonic suite and Cape St. John Group. Three mines have operated in this area including two Cu volcanogenic massive-sulphide deposits in the Betts Cove Complex (Tilt Cove and Betts Cove mines), and gold in iron-formation in the Snooks Arm Group (Nugget Pond mine). Four phases of regional deformation affected this area, including: D₁ restricted to a northwest-directed reverse fault in the Betts Cove Complex related to ophiolite obduction; D₂ regional, penetrative deformation accompanied by greenschist metamorphism; D₃ related to open upright cross folds; and D₄ in the north, related to recumbent folding coeval with extensional faults, accompanied by an amphibolitic metamorphic overprint.

RÉSUMÉ

Le sous-sol de la partie nord-est de la péninsule Baie Verte (Terre-Neuve-et-Labrador, SNRC 2-E/13 et des parties de SNRC 12-I/4, 2-E/14 et 2-E/12) est constitué du Groupe de Ming's Bight du Néoprotérozoïque à l'Ordovicien, des ophiolites du Complexe de Betts Cove du Cambrien, de la succession sous-marine de couverture des ophiolites du Groupe de Snooks Arm de l'Ordovicien, ainsi que d'un assemblage de recouvrement continental incluant la suite plutonique de Burlington de l'Ordovicien-Silurien, la suite intrusive de Stocking Harbour, la suite plutonique de Confusion Bay et le Groupe de Cape St. John, tous du Silurien. Trois mines ont été exploitées dans cette région dont deux gisements de sulfures massifs volcanogènes riches en cuivre, dans le Complexe de Betts Cove (mines Tilt Cove et Betts Cove), et un gisement d'or encaissé dans des formations de fer, dans le Groupe de Snooks Arm (mine Nugget Pond). Quatre phases de déformation régionales ont touché la région, dont : D₁, limitée à une faille inverse à vergence nord-ouest dans le Complexe de Betts Cove et reliée à l'obduction des ophiolites; D₂, une déformation pénétrative régionale accompagnée d'un métamorphisme du faciès des schistes verts; D₃, reliée à des plis transversaux, ouverts et droits; et D₄, dans la partie nord, associée à des plis couchés et à des failles d'extension contemporaines et accompagnée d'une surimpression métamorphique du faciès des amphibolites.

ABOUT THE MAP

General Information

Authors: T. Skulski, S. Castonguay, Y. Moussallam, V.J. McNicoll, C.R. van Staal, and J.H. Bédard

Geological field work by T. Skulski, S. Castonguay, Y. Moussallam (University of Ottawa), C.R. van Staal, I. Kerr (University of Victoria), and S. Hinchey (Memorial University), 2006–2008

Geological compilation and notes by T. Skulski and S. Castonguay, 2008–2012

Geophysical and remotely sensed data processed by H. Slavinski, B. Spicer, W. Morris, H. Ugalde (McMaster University), 2007–2008, G. Kilfoil (Geological Survey of Newfoundland and Labrador), 2007, and T. Skulski, 2008–2013

Compiled structural data digitized by M. Currie and A. Magee

Cartography by N. Côté

Critical review by A. Zagorevski (Geological Survey of Canada) and H.A. Sandeman (Geological Survey of Newfoundland and Labrador)

Scientific editing by E. Inglis

Initiative of the Geological Survey of Canada, conducted under the auspices of the Appalachian TGI-3 Project as part of Natural Resources Canada's Targeted Geoscience Initiative (TGI-3).

Map projection Universal Transverse Mercator, zone 21.
North American Datum 1983

Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications. Elevations above mean sea level are expressed in metres (NTS 2-E/13) and feet (NTS 2-E/12, 2-E/14, 2-L/4)

Magnetic declination 2015, 19°51'E, decreasing 11.6' annually.

This map is not to be used for navigational purposes.

Title photograph: Recumbent F₄ fold of Cape St. John Group. Confusion Bay, Newfoundland and Labrador. Photograph by S. Castonguay. 2014-111

The Geological Survey of Canada welcomes corrections or additional information from users.

Data may include additional observations not portrayed on this map.
See documentation accompanying the data.

This publication is available for free download through GEOSCAN (<http://geoscan.nrcan.gc.ca/>).

Map Viewing Files

The published map is distributed as a Portable Document File (PDF), and may contain a subset of the overall geological data for legibility reasons at the publication scale.

ABOUT THE GEOLOGY

Descriptive Notes

The Nippers Harbour map area in the northeast part of Baie Verte Peninsula (Newfoundland and Labrador) is one of three new 1:50 000 scale maps in this area that are based on new and compiled bedrock geological (including geochronological and lithogeochemical data) and remotely sensed data (Skulski et al., 2015a, b). This part of the Newfoundland Appalachians is underlain by five tectonostratigraphic entities including: Neoproterozoic to Early Paleozoic rocks of the Laurentian continental margin (Ming's Bight Group); Cambrian ophiolite (Betts Cove and Pacquet complexes); Lower Ordovician ophiolite cover (Snooks Arm Group); and Upper Ordovician to Silurian continental, volcano-plutonic rocks (Burlington plutonic suite, Confusion Bay

plutonic suite and Cape St. John Group). The IUGS time scale of Cohen et al. (2013) is utilized herein.

TECTONOSTRATIGRAPHY AND SETTING

The Ming's Bight Group (unit nPOMB) is part of the Fleur de Lys Supergroup and correlates with the Rattling Brook Group to the west (Hibbard, 1983). It comprises pelite, semipelite, psammite, and minor metaconglomerate with small pods of mafic amphibolite and fuchsite actinolite schist. The metaclastic rocks are associated with larger panels of massive to pillowed, tholeiitic metabasalt and amphibolite, termed Pelly Point schist by Hibbard (1983) and these are herein correlated with the ca. 560 Ma Birchy Complex (unit nPBm; van Staal et al., 2013). Originally described as a continental margin promontory (Hibbard, 1983), the Ming's Bight Group has since been recognized as an extensional core complex that exposes the Laurentian continental margin (Anderson et al., 2001).

The Cambrian Betts Cove Complex represents the best preserved and most complete section of ophiolite crust on Baie Verte Peninsula (Bédard et al., 2000b). The Pacquet complex to the west comprises correlative dismembered ophiolite. A sheared serpentinite (unit €Ms) at the structural base of the Betts Cove Complex is overlain and in fault contact with intrusive rocks including layered ultramafic cumulate rocks (unit €BHu), cumulate gabbro and pyroxenite (unit €BHg), massive gabbro (unit €BHM), late intrusive gabbro (unit €BHlg) dated at 488.6 ± 3.1 –1.8 Ma (location 22, Table 1), and sheeted boninite dykes (unit €BHs). The contact with the volcanic section is typically transitional and characterized by mixed sheeted dykes and pillowed boninite (unit €BHbs). The Betts Head Formation (unit €BHb), in the lower volcanic section includes sparsely porphyritic and spherulitic boninite pillow lavas and breccia units that are chemically subdivided into low- and intermediate-Ti varieties (Bédard et al., 2000b, and references therein). In contrast to the stratigraphy defined by Bédard et al. (2000b), the present authors include the volcanic rocks of the Mount Misery Formation (unit €MMm) in the Betts Cove Complex, rather than in the Snooks Arm Group (Skulski et al., 2010). The Mount Misery Formation is composed of plagioclase-phyric, pillowed island-arc tholeiitic basalt flows and breccia units that are interbedded with, and chemically transitional to the intermediate-Ti boninite units.

The Lower to Middle Ordovician Snooks Arm Group (Bédard et al., 2000b; Skulski et al., 2010) overlies the Betts Cove and Pacquet complexes. The Scrape Point Formation rests paraconformably to locally disconformably on underlying ophiolite crust. The basal Nugget Pond member (unit OSPI) comprises a local, basal sedimentary breccia with clasts of pillow basalt and chert cemented by quartz and magnetite (Sangster et al., 2007), overlain by jasper iron-formation interbedded with mafic volcanoclastic rocks. These are covered by volcanic-derived turbiditic siltstone, wacke and crystal- to lapilli-tuff units (unit OSPs) and pillowed, tholeiitic basalt (unit OSPm) and related volcanoclastic rocks (unit OSPv). The overlying Bobby Cove Formation includes a lower calc-alkaline pillow basalt and andesite (unit OBCm), overlain by a regionally distinctive marker unit (Prairie Hat member; unit OBcc) consisting of clinopyroxene-phyric to megacrystic, intermediate-, calc-alkaline crystal tuff to tuff breccia. The Prairie Hat member is overlain by a 470 ± 4 Ma (location 16, Table 1) felsic crystal tuff (unit OBCf), and an upper sequence of mafic volcanoclastic (unit OBCv) and turbiditic, mafic

epiclastic rocks (unit OBCE). An iron-formation marker unit (unit OBCi) lies at the top of the Bobby Cove Formation and consist of banded jasper or quartz-magnetite iron-formation. The Venams Bight Formation consists of tholeiitic, pillowed basalt and related breccia (unit OVBm). The overlying Balsam Bud Cove Formation consists of black, graptolitic shale and turbiditic wacke (unit OBBs) covered by a felsic volcanic unit (unit OBBf) including a 467 ± 4 Ma (location 14, Table 1) rhyolite overlying the Betts Cove Complex, and 470 ± 4 Ma (location 15, Table 1) lapilli tuff overlying the Pacquet complex. Thick-bedded, volcanic-derived conglomerate forms the upper part of the formation at Betts Cove. The Round Harbour Formation marks the top of the Snooks Arm Group and consists of pillowed tholeiitic basalt and volcanogenic sedimentary rocks (unit ORHm). Numerous synvolcanic, tholeiitic gabbro sills and dykes intrude the Snooks Arm Group (unit OSAg) and boninite units of the Pacquet complex.

Silurian (Llandoveryan) plutonic rocks of the Burlington plutonic suite intrude the ophiolite complexes and the Snooks Arm Group cover. These include calc-alkaline, hornblende-biotite quartz diorite (unit SBqd) and biotite±hornblende granodiorite (unit SBgd) dated at 441 ± 1.2 Ma (location 9, see Skulski et al., 2015b). The Stocking Harbour intrusive suite (unit SSHg) intrudes the Burlington plutonic suite and includes peraluminous, biotite±muscovite tonalite-granodiorite-trondhjemite and muscovite-garnet leucogranodiorite (Epstein, 1983). The Wenlockian to Ludlovian Confusion Bay plutonic suite (Cape Brulé Porphyry, Dunamagon granite, Reddits Cove gabbro, Seal Island Bight syenite, and La Scie granite) and Cape St. John Group are coeval and share common tholeiitic to alkaline parental magmas. The Cape Brulé Porphyry is a composite early subvolcanic pluton comprising tholeiitic, two feldspar–biotite–quartz-phyric monzogranite (unit SCBp) dated at 429 ± 4 Ma (location 6, Table 1), with a melanocratic, coarse-grained, granodiorite to monzogranite phase (unit SCBi), intrusive breccia (unit SCBix), and younger quartz-feldspar porphyry-aplite ring dyke (unit SCBd). An angular unconformity separates the Cape St. John Group from the underlying Snooks Arm Group, Burlington plutonic suite, and Pacquet complex. A lenticular lower sedimentary sequence (Beaver Cove formation; unit SBCs) comprises red-buff-weathering arkose and conglomerate. This is overlain by amygdaloidal tholeiitic basalt, alkaline basalt and hawaiite flows of the Long Pond formation (unit SLPm). These are covered by welded rhyolitic tuff of the Goat Pond formation (unit SGPw) dated at 427.8 ± 0.6 Ma (location 5, Table 1). Overlying these are rhyolitic tuff breccia with jasper and fuchsite clasts in a crystal-tuff matrix of the Red Cliff Pond formation (unit SRCpb) and a younger rhyolitic tuff breccia with blocks of basalt, rhyolite, and felsic tuff in a mafic, highly magnetic matrix (Goss Pond formation; unit SGPb). These are in turn overlain by fine-grained rhyolitic to comenditic tuff (Binch Pond formation; unit SBpt) and amygdaloidal basalt, alkaline basalt, hawaiite and transitional basalt of the Mathis Pond formation (unit SMPm). Overlying intermediate to felsic lapilli tuff units are subdivided into the Armchair Pond formation (unit SAPI) containing black lapilli, bombs, and recrystallized fiamme; and the Little Caplin Cove formation (unit SLCl) with beige lapilli. These are overlain by a 426.1 ± 0.5 Ma (location 2, Table 1) flow-banded rhyolite that includes K-feldspar–phyric rhyolite (Welsh's Pond formation; unit SWPr) that grades westward into tuff breccia, and overlying, comparatively aphyric flow-banded, high-K rhyolite of the Blanche Point formation (unit SBPr). A welded, high-K, quartz–two-

feldspar–magnetite felsic tuff (Carolina Point formation; unit SCPr) is overlain by a white, high-K, K-feldspar porphyritic lapilli tuff with elongate sericitized pumice lapilli (Brent's Cove formation; unit SBCL). These are overlain by a thin, amygdaloidal aphyric basalt unit (Stakes Pond formation; unit SSPm) and an upper flow-banded aphyric rhyolite (La Scie formation; unit SLSr). Younger phases of the Confusion Bay plutonic suite include the medium- to coarse-grained Dunamagon monzogranite (unit SDgr) dated at 427.2 ± 1.4 Ma (location 4, Table 1), equigranular to layered Reddits Cove gabbro (unit SRg), fine- to medium-grained, Seal Island Bight syenite (unit SSIB) dated at 426.3 ± 1.1 Ma (location 3, Table 1), and LaScie granite (unit SLSG) locally with abundant mafic enclaves (unit SLGa).

Two caldera-collapse cycles are recognized in the Cape St. John Group and related Confusion Bay plutonic suite (this study). The Cape Brulé Porphyry is interpreted as a subvolcanic, caldera-related granite. Evidence for high-level emplacement include prominent quartz plus two-feldspar porphyritic character in a fine-grained groundmass, presence of miarolitic cavities, reverse compositional zoning with a melanocratic core (unit SCBix) and leucocratic rim (unit SCBp), roof pendants of ophiolitic crust, dykes and plugs of intrusive felsic breccia (unit SCBix), and a half-ring dyke (unit SCBd) of granite porphyry. The extrusive equivalent of the Cape Brulé Porphyry (429 ± 4 Ma; location 6, Table 1) is the precollapse welded tuff of the Goats Pond formation (unit SGPw; 427.8 ± 0.6 Ma; location 5, Table 1). The first cycle of caldera collapse may be marked by deposition of volcanic breccia units; initially with abundant ophiolite and cover-related basement xenoliths (unit SRCPb) and later with cognate xenoliths (unit SGPb) culminating with caldera-fill tuff (unit SBPt). Resurgence of the caldera resulted in a ring dyke forming in a trap-door caldera uplift (unit SCBd) followed by eruption of mildly peralkaline rhyolitic tuff (unit SBPt). A period of intracaldera resurgence involved eruption of lapilli tuff units (units SAPI and SLCL), followed by eruption of flow-banded rhyolite domes (units SWPr at 426.1 ± 0.5 Ma; location 2, Table 1, and SBPr). The second caldera collapse event may be distal (King's Point caldera at 427 ± 2 Ma; Coyle, 1990) and involved eruption of high-K welded ash-flow tuff (unit SCPr), eruption of high-K lapilli tuff (unit SBCL), and followed by later resurgence and eruption of rhyolite domes (unit SLSr).

ECONOMIC GEOLOGY

Three mines have operated in the Nippers Harbour map area (Sangster et al., 2007, and references therein). Copper-rich, ophiolite-hosted volcanogenic massive sulphide was mined in the Tilt Cove area, where ore lenses were hosted by the Betts Head and Mount Misery formations. The Betts Cove deposit comprises copper-rich massive sulphide hosted by Betts Head formation boninite and underlying sheeted dykes. The Nugget Pond mine is a past-producing gold mine where mineralization was hosted by quartz-albite-carbonate alteration in iron-formation in the Nugget Pond member of the Scrape Point Formation (Snooks Arm Group).

TECTONOMETAMORPHIC EVOLUTION

At least four phases of regional deformation have affected the rocks of the eastern Baie Verte Peninsula (DeGrace et al., 1976; Hibbard, 1983; Stella, 1987; Tremblay et al., 1997; Anderson et al., 2001; Castonguay et al., 2009; Skulski et al., 2010, among

others). In general, rocks show a general increase in strain and metamorphic grade from south to north across the map area.

Structures related to D_1 (D_e of Hibbard, 1983) are weakly developed or poorly preserved. In the Betts Cove Complex, a northwest-directed reverse fault, interpreted as a D_1 obduction-related structure (Tremblay et al., 1997; Bédard et al., 2000b), juxtaposes the Betts Head and Mount Misery formations against cumulate peridotite, sheeted dykes, and serpentinite. The D_1 phase is interpreted as related to obduction of ophiolite rocks, arc collision of the Baie Verte Oceanic Tract, and attempted subduction of the autochthonous margin during the Taconic Orogeny (Waldron et al., 1998; van Staal et al., 2007). Post- D_1 , pre- D_2 north- to northwest-dipping brittle-ductile normal faults were interpreted by Tremblay et al. (1997) to mark most of the contact between the northern Betts Cove Complex and the Cape St. John Group, and southward, between the talc-serpentine schist and the Snooks Arm Group or Cape Brulé Porphyry.

Penetrative D_2 deformation (D_m of Hibbard (1983); D_1 of Anderson et al. (2001)) is accompanied by greenschist-facies metamorphism. In the southern part of the peninsula, S_2 is a steep to moderately northeast- to northwest-dipping spaced cleavage. Mesoscopic F_2 folds are rarely observed, but reversal of pillow tops and map-scale distribution of marker horizons suggest that macroscopic F_2 folds are present. In the northern part, the S_2 schistosity is axial planar to a series of major close to tight, upright to overturned east- to northeast-trending F_2 folds. The D_2 fabrics are associated with ductile to brittle-ductile south- to southeast-directed shear zones, such as the fault that juxtaposes the Cape St. John Group and Cape Brulé Porphyry against the Betts Cove Complex (Tremblay et al., 1997; Bédard et al., 2000b). The D_3 deformation is characterized by open, upright F_3 cross folds that vary in style and fold plunge depending on the orientation of earlier structures. These north-trending folds are locally associated with a subvertical axial-planar cleavage. The D_2 and D_3 structural phases are interpreted to form part of a progressive deformation associated with the Salinic Orogeny (Skulski et al., 2010) that led to strain partitioning in an overall sinistral transpressional regime (Waldron et al., 1998). Salinic deformation and metamorphism are locally constrained to be less than 426 Ma (location 2, Table 1; age of D_2 deformed upper Cape St. John Group) and as young as 417 Ma (location 41 and 103, Table 1; $^{40}\text{Ar}/^{39}\text{Ar}$ hornblende and biotite ages). The later age (417 Ma) constrains the age of metamorphism of the ca. 440 Ma phase of the Burlington plutonic suite (units SBgd and SBqd), insofar as it is a depositional basement to the ca. 427 Ma King's Point volcanic complex in southeastern Baie Verte Peninsula.

The fourth deformation phase (D_L of Hibbard (1983); D_{2-3} of Anderson et al. (2001)) is recognized in the northern part of the map area. It is characterized by shallowly inclined to recumbent, open to tight F_4 folds that are interpreted as cogenetic with southeast-dipping ductile-brittle extensional shear zones (Anderson et al., 2001). An axial-planar S_4 fabric, mainly formed by transposition of S_2 , dips shallowly to the southeast. D_4 is locally the dominant structure in the Ming's Bight Group, where it is associated with an amphibolite-facies metamorphic overprint (with $^{40}\text{Ar}/^{39}\text{Ar}$ hornblende ages from 405 Ma (location 65, Table 1) to 359 Ma (location 59, Table 1); 385 Ma titanite (Location 30, Table 1), and $^{40}\text{Ar}/^{39}\text{Ar}$ muscovite from 365 Ma (location 82, Table 1) to 358 Ma (location 84, Table 1); Anderson et al., 2001), which has also affected the northwestern exposed part of the Cape St. John Group. Later (D_3 of Anderson et al., 2001) moderately to steeply south- and north-dipping ductile-brittle normal faults overprint

contacts of the Ming's Bight Group. The D₄ phase is interpreted to have initiated progressive extensional unroofing of the Ming's Bight Group during an overall dextral strike-slip (transpressional to transtensional) regime (Waldron et al., 1998; Anderson et al., 2001). Several northeast-trending faults occur in the southwestern part of the map area. They affect the Silurian intrusions with systematic apparent dextral offsets and are probably related to the Green Bay Fault, running offshore Notre Dame Bay (Hibbard, 1983).




Figure 1. Map sources: 1 = Anderson, 1998; 2a = Bédard et al., 1999; 2b = Bédard et al., 2000b; 3 = Coates, 1970; 4 = Colman-Sadd and Crisby-Whittle; 5 = DeGrace et al., 1976; 6 = Epstein, 1983; 7 = Hibbard, 1983; 8 = Piercey, 1996; 9 = Sheppard et al., 1987; 10 = Stella, 1987.




Figure 2. a) Tectonic map of Newfoundland and b) simplified geological map of the Baie Verte Peninsula (modified from Hibbard, 1983; Skulski et al., 2010; van Staal et al., 2013).

| Location | Method | Mineral | Rock type | Code | Age (Ma) | Interpretation | Note | Reference |
|--|---|-----------|------------------------------------|-------|------------------|-------------------------|--|-----------------------------|
| 2 | U/Pb-TIMS | Zircon | Flow-banded rhyolite | SWPr | 426.1 ± 0.5 | Crystallization age | | Skulski et al., 2012 |
| 3 | U/Pb-TIMS | Zircon | Coarse-grained syenite | SSib | 426.3 ± 1.1 | Crystallization age | | Skulski et al., 2012 |
| 4 | U/Pb-TIMS | Zircon | Monzogranite | SDgr | 427.2 ± 1.4 | Crystallization age | | Skulski et al., 2012 |
| 5 | U/Pb-TIMS | Zircon | Welded tuff | SGPw | 427.8 ± 0.6 | Crystallization age | | Skulski et al., 2012 |
| 6 | U/Pb SHRIMP | Zircon | Quartz-feldspar monzogranite | SCBp | 429 ± 4 | Crystallization age | | Skulski et al., 2012 |
| 14 | U/Pb SHRIMP | Zircon | Rhyolite flow | OBBf | 467 ± 4 | Crystallization age | | Skulski et al., 2010 |
| 15 | U/Pb SHRIMP | Zircon | Felsic lapilli tuff | OBBf | 470 ± 4 | Crystallization age | | Skulski et al., 2010 |
| 16 | U/Pb SHRIMP | Zircon | Felsic tuff | OBcc | 470 ± 4 | Crystallization age | | Skulski et al., 2010 |
| 22 | U/Pb-TIMS | Zircon | Coarse-grained quartz gabbro | CBHg | 488.6 ± 3.1/-1.8 | Crystallization age | | Dunning and Krogh, 1985 |
| 30 | U/Pb-TIMS | Titanite | Biotite monzogranite | SDgr | 385 ± 1 | Metamorphism | | Skulski et al., 2012 |
| 32 | U/Pb-TIMS | Rutile | Chlorite schist (shear zone) | nPOMB | 380 ± 2 | Cooling age | | Anderson et al., 2001 |
| 33 | U/Pb-TIMS | Microlite | Pegmatite cuts F ₃ fold | nPOMB | 355 ± 2 | Cooling age | | Anderson et al., 2001 |
| 40 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Granodiorite | SBgd | 421 ± 5 | Metamorphic cooling age | See note 3. | Dallmeyer and Hibbard, 1984 |
| 41 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Granodiorite | SBgd | 417 ± 5 | Metamorphic cooling age | See note 3. | Dallmeyer and Hibbard, 1984 |
| 44 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Schist | OVBm | 357 ± 5 | Metamorphic cooling age | See note 3. | Dallmeyer and Hibbard, 1984 |
| 46 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Schist | nPBm | 369 ± 5 | Metamorphic cooling age | Recalculated as 4 step plateau (98% ³⁹ Ar); see note 2. | Dallmeyer, 1977 |
| 48 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Schist | OVBm | 352 ± 5 | Metamorphic cooling age | See note 3. | Dallmeyer and Hibbard, 1984 |
| 59 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Schist | nPBm | 359 ± 5 | Metamorphic cooling age | See note 3. | Dallmeyer and Hibbard, 1984 |
| 60 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Mafic dyke | SDgr | 379 ± 2 | Metamorphic cooling age | See note 3. | Anderson et al., 2001 |
| 61 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Mafic dyke | nPOMB | 382 ± 8 | Metamorphic cooling age | See note 3. | Anderson et al., 2001 |
| 62 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Mafic dyke | nPOMB | 390 ± 9 | Metamorphic cooling age | Fault-bounded mafic sliver; see note 3. | Anderson et al., 2001 |
| 63 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Gabbro | OSu | 477 ± 6 | | See note 3. | Anderson et al., 2001 |
| 64 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Amphibolite | OSu | 379 ± 21 | Metamorphic cooling age | Amphibolite mylonite, fault sliver; see note 3. | Anderson et al., 2001 |
| 65 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Amphibolite | OBce | 405 ± 8 | Metamorphic cooling age | Polydeformed amphibolite; see note 3. | Anderson et al., 2001 |
| 66 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Amphibole | Amphibolite | OBBs | 384 ± 5 | Metamorphic cooling age | Mylonitic amphibolite; see note 3. | Anderson et al., 2001 |
| 78 | ⁴⁰ Ar/ ³⁹ Ar laser step-heating | Muscovite | Schist | SMPm | 379.9 ± 2.1 | Metamorphic cooling age | Pseudo-plateau, 81% gas released, 1 step dropped. | Castonguay et al., 2010 |
| 81 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Pegmatite | nPOMB | 365 ± 3 | Metamorphic cooling age | See note 4. | Anderson et al., 2001 |
| 82 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Schist | nPOMB | 365 ± 3 | Metamorphic cooling age | Muscovite defines S ₁ ; see note 4. | Anderson et al., 2001 |
| 83 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Schist | nPOMB | 361 ± 3 | Metamorphic cooling age | Big Brook shear zone; see note 4. | Anderson et al., 2001 |
| 84 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Pegmatite | nPOMB | 358 ± 3 | Metamorphic cooling age | Deformed pegmatite dyke; see note 4. | Anderson et al., 2001 |
| 85 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Schist | nPOMB | 364 ± 3 | Metamorphic cooling age | Muscovite defines S ₁ ; see note 4. | Anderson et al., 2001 |
| 86 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Schist | nPOMB | 362 ± 3 | Metamorphic cooling age | Muscovite defines S ₁ ; see note 4. | Anderson et al., 2001 |
| 87 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Schist | nPOMB | 361 ± 3 | Metamorphic cooling age | See note 4. | Anderson et al., 2001 |
| 88 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Pegmatite | nPOMB | 364 ± 2 | Metamorphic cooling age | Deformed pegmatite; see note 4. | Anderson et al., 2001 |
| 89 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Muscovite | Schist | nPOMB | 366 ± 3 | Metamorphic cooling age | Muscovite defines S ₁ ; see note 4. | Anderson et al., 2001 |
| 91 | ⁴⁰ Ar/ ³⁹ Ar laser step-heating | Muscovite | Felsic phyllite | OBBf | 369.3 ± 2.0 | Metamorphic cooling age | Combined plateau from two aliquots, 86% gas released. | Castonguay et al., 2010 |
| 93 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Biotite | Schist | OVBm | 342 ± 5 | Metamorphic cooling age | See note 3. | Dallmeyer and Hibbard, 1984 |
| 103 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Biotite | Granodiorite | SBgd | 417 ± 10 | Metamorphic cooling age | Discordant spectrum; reported plateau age (6 steps); see note 3. | Dallmeyer and Hibbard, 1984 |
| 104 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Biotite | Schist | nPOMB | 352 ± 5 | Metamorphic cooling age | Total gas age; see note 2. | Dallmeyer, 1977 |
| 105 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Biotite | Schist | nPOMB | 339 ± 4 | Metamorphic cooling age | Four-step plateau age; see note 3. | Dallmeyer and Hibbard, 1984 |
| 106 | ⁴⁰ Ar/ ³⁹ Ar furnace step-heating | Biotite | Granite | SDgr | 346 ± 5 | Metamorphic cooling age | Reported plateau age (5 step); see note 3. | Dallmeyer and Hibbard, 1984 |
| 108 | ⁴⁰ Ar/ ³⁹ Ar laser step-heating | Biotite | Schist | SMPm | 378.6 ± 2.1 | Metamorphic cooling age | Pseudo-plateau, 35% gas released. | Castonguay et al., 2010 |
| Notes 1 ⁴⁰ Ar/ ³⁹ Ar ages (new and historical data) have been calculated using a total ⁴⁰ K decay constant of 5.463E-10 (Min et al., 2000). Some plateau ages (indicated in table) have been recalculated using IsoPlot version 3.7 of Ken Ludwig using recalculated ages of steps (in light of revised decay constant and internal standard ages) using Noah Mclean's ArArReCalc, 7-31-09. 2 Recalculated using age of Biotite standard SB-2 as 164 Ma revised from original value of 160.2 Ma in light of revised ⁴⁰ K decay constant (note 1). 3 Recalculated using an age of MMhb-1 hornblende standard of 523.1 Ma (Renne et al., 1998) revised from original data calculated using an age of MMhb-1 of 519.5 Ma (Alexander et al., 1978). 4 Recalculated using an age of MMhb-1 hornblende standard of 523.1 Ma (Renne et al., 1998) revised from original data calculated using an age of MMhb-1 of 520 Ma (Samson and Alexander, 1987). | | | | | | | | |

Table 1. Geochronological data.

| Location | Name | Status | Commodity | Secondary commodity |
|----------|------------------|---------------|-----------|----------------------------|
| 1 | Betts Cove mine | Past producer | Copper | Gold, zinc |
| 2 | Tilt Cove mine | Past producer | Copper | Gold, nickel, silver, zinc |
| 3 | Nugget Pond mine | Past producer | Gold | Copper, silver |

Table 2. Past-producing mines.

| Location | Name | Commodity | Secondary commodity |
|----------|------------------------|-----------|------------------------------------|
| 6 | Footie Pond | Copper | Zinc, gold, silver |
| 7 | Rogues Harbour | Copper | Gold, lead |
| 13 | Nippers Harbour Copper | Copper | Iron |
| 27 | Mount Misery | Copper | Gold, copper, zinc, silver |
| 40 | Long Pond (Nudulama) | Copper | Gold, lead, iron, |
| 41 | Muir's Pond | Copper | Silver, gold, iron, nickel, cobalt |
| 44 | Mud Pond | Gold | Copper, nickel, talc |

Table 3. Drilled prospects.

References

Alexander, E.C., Jr., Mickelson, G.M., and Lanphere, M.A., 1978. MMhb-1; a new ^{40}Ar - ^{39}Ar dating standard; in 4th International Conference on Geochronology, Cosmochronology, and Isotope Geology, Snowmass-at-Aspen, Colorado, 1978. Short Papers: U.S. Geological Survey, Open-File Report 78-701, p. 6–8.

Anderson, S.D., 1998. Structure, metamorphism, and U–Pb and $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of the Ming's Bight Group, and the Paleozoic tectonic evolution of the Baie Verte Peninsula, Newfoundland; Ph.D. thesis, Dalhousie University, Halifax, Nova Scotia, 452 p.

Anderson, S.D., Jamieson, R.A., and Reynolds, P.H., 2001. Devonian extension in northwestern Newfoundland: $^{40}\text{Ar}/^{39}\text{Ar}$ and U–Pb data from the Ming's Bight area, Baie Verte Peninsula; *Journal of Geology*, v. 10, p. 191–211.

Bédard, J.H., Lauzière, K., Sangster, A., Tellier, M., Tremblay, A., and Dec, T., 1999. Geology, Betts Cove Ophiolite and its cover rocks, Newfoundland; Geological Survey of Canada, Map 1969A, scale 1:20 000. doi:10.4095/211851

Bédard, J.H., Lauzière, K., Boisvert, É., Deblonde, C., Sangster, A., Tremblay, A., and Dec, T., 2000a. Betts Cove geological dataset for GIS applications; Geological Survey of Canada, Open File 3623, 1 CD-ROM. doi:10.4095/211544

Bédard, J.H., Lauzière, K., Tremblay, A., Sangster, A., Douma, S.L., and Dec, T., 2000b. Betts Cove ophiolite and its cover rocks, Newfoundland; Geological Survey of Canada, Bulletin 550, 76 p. doi:10.4095/211642

Castonguay, S., Skulski, T., van Staal, C., and Currie, M., 2009. New insights on the structural geology of the Pacquet Harbour Group and Point Rousse Complex, Baie Verte Peninsula, Newfoundland; Newfoundland and Labrador Department of Natural Resources, Geological Survey, Current Research 2009, Report 09-1, p. 147–158.

Castonguay, S., Skulski, T., van Staal, C.R., McNicoll, V., and Joyce, N., 2010. Revisiting the Baie Verte Flexure: from Silurian transpression to Devonian transtension, a long-lived oblique transfer zone, Baie Verte Peninsula, Newfoundland Appalachians; *in* Abstracts with Programs, Geological Society of America, Northeastern and Southeastern Sections Joint Annual Meeting, March 13–16, Baltimore, Maryland, p. 164.

Coates, H.J., 1970. Structural and metamorphic history of the Pacquet Harbour-Grand Cove area, Burlington Peninsula, Newfoundland; M.Sc. thesis, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador, 79 p.

Cohen, K.M., Finney, S.M., Gibbard, P.L., and Fan, J.-X., 2013. The ICS International Chronostratigraphic Chart; *Episodes*, v. 36, p. 199–204.

Colman-Sadd, S.P. and Crisby-Whittle, L.V.J., 2002. Partial bedrock geology dataset of the Island of Newfoundland (NTS areas 02E, 12H, 12G, and parts of 01M, 02D, 02L, 12A, 12B, and 12I); Newfoundland and Labrador Department of Mines and Energy, Open File NFLD/2616.

Coyle, M., 1990. Geology, geochemistry and geochronology of the Springdale Group, an Early Silurian caldera in central Newfoundland; Ph.D. thesis, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador, 310 p.

Dallmeyer, R.D., 1977. $^{40}\text{Ar}/^{39}\text{Ar}$ age spectra of minerals from the Fleur de Lys terrane in northwest Newfoundland: their bearing on chronology of metamorphism within the Appalachian orotectonic zone; *Journal of Geology*, v. 85, p. 89–103.

Dallmeyer, R.D., and Hibbard, J., 1984. Geochronology of the Baie Verte Peninsula, Newfoundland: implications for the tectonic evolution of the Humber and Dunnage Zones of the Appalachian Orogen; *Journal of Geology*, v. 92, p. 489–512.

DeGrace, J.R., Kean, B.F., Hsu, E., and Green, T., 1976. Geology of the Nippers Harbour area (2E/13), Newfoundland; Department of Mines and Energy, Newfoundland, Report 76-3, 73 p., scale 1:50 000.

Dunning, G.R. and Krogh, T.E., 1985. Geochronology of ophiolites of the Newfoundland Appalachians; *Canadian Journal of Earth Sciences*, v. 22, p. 1659–1670.

Epstein, R.S., 1983. The eastern margin of the Burlington granodiorite, Newfoundland; M.Sc. thesis, University of Western Ontario, London, Ontario, 188 p.

Hibbard, J., 1983. Geology of the Baie Verte Peninsula, Newfoundland; Department of Mines and Energy, Government of Newfoundland and Labrador, Memoir 2, 279 p., scale 1:100 000.

Min, K., Mundil, R., Renne, P.R., and Ludwig, K.R., 2000. A test for systematic errors in $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology through comparison with U/Pb analysis of a 1.1 Ga rhyolite; *Geochimica et Cosmochimica Acta*, v. 64, p. 73–98.

Piercey, J.P., 1996. The geology and geochemistry of the Southern Pacquet Harbour Group volcanics, Baie Verte Peninsula, Newfoundland; B.Sc. thesis, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador, 123 p.

Renne, P.R., Swisher, C.C., Deino, A.L., Karner, D.B., Owens, T.L., and Depaolo, D.J., 1998. Intercalibration of standards, absolute ages and uncertainties in $^{40}\text{Ar}/^{39}\text{Ar}$ dating; *Chemical Geology*, v. 145, p. 117–152.

Samson, S.D. and Alexander, E.C., 1987. Calibration of the interlaboratory $^{40}\text{Ar}/^{39}\text{Ar}$ dating standard, MMhb-1; *Chemical Geology, Isotope Geoscience Section*, v. 66, p. 27–34.

Sangster, A.L., Douma, S.M., and Lavigne, J., 2007. Base metal and gold deposits of the Betts Cove complex, Baie Verte Peninsula, Newfoundland; *in* Mineral Deposits of Canada: A Synthesis of Major Deposit-Types, District Metallogeny, the Evolution of Geological Provinces, and Exploration Methods, (ed.) W.D. Goodfellow; Geological Association of Canada, Mineral Deposits Division, Special Publication, no. 5, p. 703–721.

Sheppard, B., Strickland, R., Mercer, B., and Delaney, P., 1987. Exploration Report on Baie Verte Property licence# 3041, CB-5015-5017, 5020-5023, License 3154, CB 5330.b; Sharpe Energy and Resources Ltd., Newfoundland Mineral, Lands and Mines Division, Assessment Report Nfld1866, 183 p.

Skulski, T., Castonguay, S., McNicoll, V., van Staal, C.R., Kidd, W., Rogers, N., Morris, W., Ugalde, H., Slavinski, H., Spicer, W., Mousallam, Y., and Kerr, I., 2010. Tectonostratigraphy of the Baie Verte Oceanic Tract and its ophiolite cover sequence on the Baie Verte Peninsula, Newfoundland; Newfoundland and Labrador Department of Natural Resources, Geological Survey, Current Research 2010, Report 10-1, p. 315–335.

Skulski, T., McNicoll, V., Whalen, J.B., Moussallam, Y., Dunning, G., Castonguay, S., Cawood, P., Kidd, W.S.F., and van Staal, C., 2012. Subduction to slab-break-off transition recorded in the timing, composition and setting of early Silurian volcano-plutonic complexes, Baie Verte Peninsula, Newfoundland; *in* Program with Abstracts, Annual Meeting Geological Association of Canada–Mineralogical Association of Canada, St. John's, Newfoundland and Labrador, May 27–29, 2012, v. 35, p. 128.

Skulski, T., Castonguay, S., Kidd, W.S.F., McNicoll, V.J., and van Staal, C.R., 2015a. Geology, King's Point, Newfoundland and Labrador, NTS 12-H/9; Geological Survey of Canada, Canadian Geoscience Map 156, scale 1:50 000. doi:10.4095/295864

Skulski, T., Castonguay, S., Kidd, W.S.F., McNicoll, V.J., van Staal, C.R., and Hibbard, J.P., 2015b. Geology, Baie Verte and parts of Fleur de Lys, Newfoundland and Labrador, NTS 12-H/16 and part of NTS 12-I/1; Geological Survey of Canada, Canadian Geoscience Map 159, scale 1:50 000. doi:10.4095/295865

Stella, P.J., 1987. Geology of the Northern Baie Verte Peninsula, Newfoundland, Canada; M.Sc. thesis, State University of New York in Albany, Albany, New York, 107 p., scale 1:15 840.

Tremblay, A., Bédard, J.H., and Lauzière, K., 1997. Taconian obduction and Silurian exhumation of the Betts Cove ophiolite, Canadian Appalachians; *Journal of Geology*, v. 105, p. 701–716.

van Staal, C.R., Chew, D.M., Zagorevski, A., McNicoll, V., Hibbard, J., Skulski, T., Castonguay, S., Escayola, M.P., and Sylvester, P.J., 2013. Evidence of Late Ediacaran hyperextension of the Laurentian-Iapetus margin in the Birchy Complex, Baie Verte Peninsula, northwest Newfoundland: implications for the opening of Iapetus, formation of peri-Laurentian microcontinents and Taconic-Grampian orogenesis; *Geoscience Canada*, v. 40, p. 94–117.

van Staal, C.R., Whalen, J.B., McNicoll, V.J., Pehrsson, S., Lissenberg, C.J., Zagorevski, A., van Breemen, O., and Jenner, G.A., 2007. The Notre Dame arc and the Taconic orogeny in Newfoundland; *in* 4-D Framework of Continental Crust, (ed.) R.D. Hatcher, Jr., M.P. Carlson, J.H. McBride, and J.R. Martínez Catalán; Geological Society of America, Memoir 200, p. 511–552.

Waldron, J.W.F., Anderson, S.D., Cawood, P.A., Goodwin, L.B., Hall, J., Jamieson, R.A., Palmer, S.E., Stockmal, G.S., and Williams, P.F., 1998. Evolution of the Appalachian Laurentian margin: Lithoprobe results in western Newfoundland; *Canadian Journal of Earth Sciences*, v. 11, p. 1271–1287.

Author Contact

Questions, suggestions, and comments regarding the geological information contained in the data sets should be addressed to:

Tom Skulski
Geological Survey of Canada
601 Booth Street
Ottawa, Ontario
K1A 0E8
Tom.Skulski@canada.ca

Coordinate System

Projection: Universal Transverse Mercator

Units: metres

Zone: 21

Horizontal Datum: NAD83

Vertical Datum: mean sea level

Bounding Coordinates

Western longitude: 56°00'00"W

Eastern longitude: 55°27'00"W

Northern latitude: 49°43'00"N

Southern latitude: 50°03'00"N

Data Model Information

This Canadian Geoscience Map does not conform to either the Bedrock or Surficial Mapping Geodatabase Data Models. The author may have included a complete description of the feature classes and attributes in the Data\Data Model Info folder.

LICENCE AGREEMENT

View the licence agreement at <http://data.gc.ca/eng/open-government-licence-canada>

ACCORD DE LICENCE

Voir l'accord de licence à <http://donnees.gc.ca/fra/licence-du-gouvernement-ouvert-canada>