




Natural Resources  
Canada

Ressources naturelles  
Canada

መ ዘመን በ ስራው ነው እና የሚከተሉት ደንብ አለው  
በዚህ ደንብ መሆኑ የሚከተሉት ደንብ መ ዘመን በ 257S  
መ ዘመን በ ስራው ነው እና የሚከተሉት ደንብ መ ዘመን በ 2016-14S  
መ ዘመን በ ስራው ነው

## CLEARWATER FIORD (NORTH)

የቦርኩል, የዚህ



መ ዘመን  
የገዢ የሚከተሉት  
በዚህ ደንብ

ለጥቃቅና

መ ዘመን በ ስራው ነው እና የሚከተሉት ደንብ  
በዚህ ደንብ መሆኑ የሚከተሉት ደንብ መ ዘመን

2016

Canada



ΔL°Q ȐC⁹C>LQΔ⁹b⁹R⁹C⁹ ȐR⁹U⁹U⁹b⁹σ⁹σ⁹b ȐL Cσ⁹bΔbC⁹C⁹σ⁹b  
R⁹D⁹L⁹C⁹D⁹R⁹σ⁹L⁹ ȐR⁹d⁹σ⁹ ȐU⁹b⁹U⁹ ȐR⁹b⁹C⁹Y⁹L⁹J⁹ NRCan-d⁹σ⁹.  
⁹b⁹D⁹A⁹b⁹σ⁹P⁹L⁹J⁹, NRCan-d⁹d⁹ Ȑb⁹D⁹A⁹b⁹A⁹R⁹C⁹R⁹D⁹σ⁹  
[nrcan.copyrightdroitdauteur.rnccan@canada.ca](mailto:nrcan.copyrightdroitdauteur.rnccan@canada.ca).

## ȐC⁹p⁹L⁹R⁹C⁹ D⁹b⁹C⁹D⁹Y⁹L⁹C⁹ Ȑb⁹D⁹A⁹b⁹σ⁹

St-Onge, M.R., Weller, O.M., Dyck, B.J., Rayner, N.M., Chadwick T., ȐL Liikane, D., 2016. Ȑa⁹σ⁹ Ȑb⁹D⁹A⁹b⁹σ⁹, Clearwater Fiord (north), ȐR⁹b⁹C⁹, Ȑa⁹σ⁹; Ȑa⁹p⁹L⁹C⁹R⁹σ⁹L⁹ Ȑb⁹D⁹A⁹b⁹b⁹ Ȑb⁹C⁹, Ȑb⁹C⁹ Ȑa⁹σ⁹ Ȑb⁹D⁹A⁹b⁹σ⁹L⁹ Ȑa⁹p⁹L⁹A⁹ 257S (ȐR⁹d⁹σ⁹); Ȑa⁹σ⁹C⁹ ȐL⁹b⁹b⁹σ⁹ Ȑb⁹D⁹A⁹b⁹b⁹ ȐU⁹D⁹G⁹S, L⁹D⁹A⁹S⁹ Ȑa⁹p⁹L⁹A⁹ 2016-14S, ȐR⁹σ⁹S 1:100 000. doi:10.4095/298760

## ABSTRACT

This map summarizes the field observations for the Clearwater Fiord (north) map area following eight weeks of regional and targeted bedrock mapping on western Hall Peninsula. The 2015 field campaign completes a two-decade mission to update map coverage for the whole of Baffin Island south of latitude 70°N. The bedrock is dominated by a Paleoproterozoic metaplutonic suite, ranging in composition from gabbro to syenogranite, with crosscutting relations indicating a progression from mafic to silicic magmatism. Prevailing upper amphibolite to lower granulite facies metamorphic conditions overlap the stability limits of magnetite and orthopyroxene, which is consistent with equilibrium phase diagrams and regional aeromagnetic data. Metasedimentary rocks, including quartzite, pelite, marble, and metagreywacke, are present as screens and enclaves between and within plutonic bodies. An examination of the 'ghost' stratigraphy suggests that the metasedimentary rocks can be correlated with the middle Paleoproterozoic Lake Harbour Group in the south and Piling Group in the north. Two basaltic dyke swarms and shallowly dipping Ordovician limestone respectively crosscut and overly the Paleoproterozoic units.

## ȐR⁹L⁹R⁹L⁹R⁹C⁹

Ȑa⁹Q Ȑa⁹p⁹L⁹A⁹ Ȑa⁹p⁹L⁹Δ⁹C⁹ Ȑb⁹D⁹A⁹b⁹R⁹C⁹ Ȑb⁹D⁹L⁹R⁹C⁹ Clearwater Fiord (north)  
Ȑa⁹p⁹L⁹A⁹ 8-σ⁹ Ȑa⁹p⁹L⁹R⁹Y⁹σ⁹ ȐR⁹C⁹b⁹Y⁹L⁹T⁹C⁹ ȐL C⁹ Ȑb⁹D⁹A⁹b⁹C⁹A⁹C⁹L⁹C⁹C⁹  
Ȑp⁹A⁹U⁹Δ⁹a⁹C⁹C⁹ Ȑa⁹p⁹L⁹C⁹D⁹σ⁹T⁹C⁹A⁹S⁹a⁹S⁹ ȐR⁹T⁹C⁹. 2015-Γ Ȑb⁹D⁹A⁹b⁹σ⁹D⁹  
ȐL⁹b⁹Y⁹C⁹D⁹C⁹ Ȑa⁹d⁹L⁹σ⁹ ȐR⁹D⁹σ⁹ Ȑa⁹p⁹L⁹C⁹ Ȑp⁹A⁹U⁹P⁹R⁹ ȐC⁹b⁹G⁹j⁹S⁹U⁹C⁹C⁹C⁹  
Ȑa⁹p⁹A⁹R⁹b⁹C⁹C⁹ ȐR⁹C⁹L⁹S⁹ 70°N. Ȑp⁹A⁹U⁹Δ⁹a⁹C⁹C⁹ Ȑa⁹C⁹b⁹σ⁹U⁹S⁹  
Paleoproterozoic metaplutonic suite, Ȑi⁹A⁹r⁹b⁹σ⁹b⁹C⁹C⁹ Ȑa⁹C⁹a⁹C⁹ ȐR⁹d⁹C⁹σ⁹ gabbro  
ȐR⁹C⁹ syenogranite, Ȑp⁹C⁹r⁹b⁹C⁹D⁹U⁹C⁹ Ȑb⁹p⁹D⁹b⁹C⁹D⁹U⁹σ⁹R⁹C⁹a⁹a⁹A⁹b⁹Y⁹  
ȐR⁹C⁹a⁹σ⁹S⁹ ȐR⁹C⁹ silicic magmatism. Ȑa⁹p⁹L⁹C⁹ C⁹i⁹H⁹D⁹ Ȑp⁹D⁹U⁹b⁹S⁹  
amphibolite ȐR⁹p⁹σ⁹S⁹C⁹ granulite facies metamorphic Ȑb⁹p⁹D⁹a⁹σ⁹U⁹R⁹C⁹  
Ȑp⁹C⁹r⁹b⁹C⁹D⁹U⁹C⁹ L⁹P⁹C⁹U⁹A⁹p⁹a⁹σ⁹S⁹ ȐR⁹C⁹a⁹b⁹U⁹C⁹D⁹ magnetite ȐL C⁹ orthopyroxene,  
ȐC⁹a⁹σ⁹b⁹C⁹C⁹ ȐC⁹p⁹U⁹C⁹a⁹C⁹ ȐR⁹C⁹a⁹b⁹C⁹ ȐR⁹C⁹a⁹b⁹C⁹ ȐL C⁹ ȐR⁹C⁹b⁹Y⁹L⁹T⁹C⁹b⁹S⁹C⁹j⁹d⁹C⁹  
σ⁹A⁹b⁹b⁹C⁹b⁹σ⁹S⁹C⁹ ȐR⁹C⁹a⁹b⁹C⁹. Metasedimentary ȐL⁹b⁹b⁹Δ⁹, ȐR⁹b⁹p⁹U⁹C⁹ ȐR⁹p⁹C⁹ quartzite,  
pelite, ȐC⁹a⁹f⁹L⁹b⁹, ȐL C⁹ ȐR⁹C⁹b⁹ metagreywacke, C⁹i⁹H⁹D⁹U⁹C⁹D⁹ ȐL ȐR⁹C⁹b⁹Y⁹L⁹σ⁹C⁹

▷ດ້ວຍການ ດັບກຳມະນຸດ plutonic Δົກຕິດີ. ນິບັດຊາຍືດີ C>σນິລີສ ‘C&gt;ΔC’ ມັດຕະນູສົງລົງ  
ແກ່ອຸດັບຢັ້ງເປົ້າ metasedimentary ແລ້ວບັດ c ບົນຍຸສົງບົບວຸດົງC ສົມບັນຫຼືC  
Paleoproterozoic ຮູ່ກົງກູ່ ລັດວັດີ ສົມຍຸສົງ ດັລ >ຈົນໄດ້ວຸດົງສ. ລົງທຶນ basaltic  
ແລ້ວບັດ c ດັກຢັ້ງ ດັບລັດບັດຈຳຈັດC Ordovician limestone ນິດົດ່ວົນC>ນິດົດ ດັລ ນິດົດ  
Paleoproterozoic ບົນຍຸສົງສົງ.

## ໝວດ ໂພນ ກິ່າມ

### ຄຽງແກ່ວົນດັບ

ົບນິດົດ: M.R. St-Onge, O.M. Weller, B.J. Dyck, N.M. Rayner, T. Chadwick, ໄ ໜ.  
D. Liikane

ມັດສ ນິບັດຊາຍືດີ M.R. St-Onge, O.M. Weller, B.J. Dyck, N.M. Rayner, T. Chadwick, ໄ ໜ.  
D. Liikane, ມັດໝັ້ນດັບນິດົດ ບັດC; S. Noble-Nowdluk, T. Milton, ໄ ໜ.  
T. Rowe, ມັດໂດ ສະລັບດີ, 2015

ມັດກ ນິບັດຊາຍືດີ ອົບນິດົດ ພຣະມະນຸດ ມັດໝັ້ນດັບນິດົດ ແລ້ວ  
M.R. St-Onge ໄ ໜ. O.M. Weller, 2015

ມັດກ ນິບັດຊາຍືດີ Lc ລົງ ສົມ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ  
ຄຽງແກ່ວົນດັບ, 2.6

ມັດໝັ້ນດັບນິດົດ A. Morin, A. Ford, C. Gilbert, L. Robertson, G. Buller, ໄ ໜ. R. Buenviaje

ມັດໝັ້ນດັບນິດົດ N. Côté

ຄຽງແກ່ວົນດັບນິດົດ ນິດົດ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ  
ຄຽງແກ່ວົນດັບນິດົດ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ ດັບ.

▷ດ້ວຍການ ດັບກຳມະນຸດ Δົກຕິດີ ມັດໝັ້ນດັບນິດົດ ມັດໝັ້ນດັບນິດົດ ແລ້ວ  
ມັດໝັ້ນດັບນິດົດ ດັບ ແລ້ວມັດໝັ້ນດັບນິດົດ (GEM) ລັດວັດັບກຳກົງ  
ຄຽງແກ່ວົນດັບນິດົດ ມັດໝັ້ນດັບນິດົດ ນິບັດຊາຍືດີ ບັດC (GSC) ລັດວັດັບນິດົດ ບັດC  
ບັດC ມັດໂດ ມັດໝັ້ນດັບນິດົດ ນິບັດຊາຍືດີ C>ນິດົດ ເປົ້າ ພຣະມະນຸດ (CNGO), ມັດໂດ  
ກົງກູ່ ບັດC ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ.

Δ້ວຍການ ດັບກຳມະນຸດ C>ນິດົດ ດັບກຳມະນຸດ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ  
ຄຽງແກ່ວົນດັບນິດົດ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ ຢຸ່ມ  
PCSP 05615

ມັດໝັ້ນດັບນິດົດ ນິດົດ ດັບກຳມະນຸດ Universal Transverse Mercator, ໄ ດັບກຳມະນຸດ 19.  
North American Datum 1983

ወደ አገልግሎት የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

“የኢትዮጵያ ማኅበር መመሪያ ስም የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

ይህ ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

ይህ ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

ይህ ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

GEOSCAN (<http://geoscan.nrcan.gc.ca/>)

በዚህ ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

## መሆኑን የሚከተሉት ሰነድ በ 1:250 000

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡

## መሆኑን የሚከተሉት ሰነድ በ 1:250 000

የሚከተሉት ሰነድ የሚከተሉት ሰነድ በ 1:250 000 የሚገኘውን ስም ነው፡፡









- 3) ሙዲስርጋብዳር ለገኖችና አስታፊሮስ ልዕሊ የየበቅጥርና Paleoproterozoic supracrustal የቦሱስ (Lake Harbour Group), ክበሩለም ተብርሃን ‘Meta Incognita microcontinent’ ስት ለSt-Onge et al. (2000a), CL<sup>b</sup>d&L የተረፈገውን ለመሆኑን crust rifted ስት the Rae የሻላጊ ማኅበርንጂ Superior craton, የሻላጊ ማኅበርንጂ CLA<sup>a</sup> መር ለC<sup>b</sup>CD<sup>a</sup>L<sup>a</sup> የሚሸፍ፡ ልዕሊ
- 4) ሙዲስርጋብዳር ለገኖችና አስታፊሮስ ልዕሊ, ምርመራለሁንና ማኅበርንጂ የሚሸፍ፡ Rae craton, ልዕሊ የየበቅጥርና Paleoproterozoic supracrustal የቦሱስ (Piling Group; Wodicka et al., 2014), ስትና ለገኖችና ለማኅበርንጂ ለማኅበርንጂ Hoare Bay Group የሻላጊ Cumberland Peninsula.

የሻላጊ ሙዲስርጋብዳር 3 ልዕሊ 4 ቀልጊዜ ማኅበርንጂ ለመሸፍ፡ Cumberland batholith, ኔዕ዗ል ለማኅበርንጂ ለመሸፍ፡ ምርመራለሁንና ለማኅበርንጂ የሚሸፍ፡ ca. 1865–1845 Ma (Whalen et al., 2010). ለማኅበርንጂ Cumberland batholith ምርመራለሁንና ለማኅበርንጂ Andean-type batholith (St-Onge et al., 2009), የሻላጊ ማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ የሚሸፍ፡ lithospheric delamination ልዕሊ ሙዲስርጋብዳር (Whalen et al., 2010). CLA<sup>a</sup> ማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ Ordovician limestone strata (Blackadar, 1967).

በዚህ ሙዲስርጋብዳር ለገኖችና ምርመራለሁንና ለማኅበርንጂ የሚሸፍ፡ ምርመራለሁንና ለማኅበርንጂ THO. ምርመራለሁንና ለማኅበርንጂ ሙዲስርጋብዳር, የሻላጊ ማኅበርንጂ 3–4 የሚሸፍ፡ ምርመራለሁንና ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ Meta Incognita microcontinent ለማኅበርንጂ Rae craton ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ ca. 1880–1865 Ma (St-Onge et al., 2006). ለማኅበርንጂ (L<sup>b</sup> ለማኅበርንጂ C<sup>b</sup> ለማኅበርንጂ) C<sup>b</sup> (St-Onge et al., 2009). ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ Piling Group, ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ, ለማኅበርንጂ Lake Harbour Group, ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ C<sup>b</sup>, ለማኅበርንጂ ለማኅበርንጂ basal orthoquartzite ልዕሊ ለማኅበርንጂ ለማኅበርንጂ iron-formation ልዕሊ metagreywacke. CL<sup>a</sup> ሙዲስርጋብዳር ለማኅበርንጂ ለማኅበርንጂ C<sup>b</sup> ለማኅበርንጂ (level 3) ለማኅበርንጂ 2015–1845 Ma, ለማኅበርንጂ ለማኅበርንጂ C<sup>b</sup> ለማኅበርንጂ C<sup>b</sup> (St-Onge et al., 2007).

የሻላጊ ማኅበርንጂ 2–3 እና ደንብ (Soper River) ሙዲስርጋብዳር ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ ለማኅበርንጂ (magmatic arc) L<sup>b</sup> Rae-Meta Incognita ለማኅበርንጂ Dunphy and Ludden, 1998), ልዕሊ





tectonostratigraphically  $\text{Nd}_{\text{I}} = 2429$ – $2690 \text{ Ma}$  Lake Harbour Group Ld $\sigma^{\text{m}}$  siliciclastic units  $\text{Nd}_{\text{I}} = 2500$ – $2700 \text{ Ma}$ .

Metagreywacke ( $\Delta_{\text{C}} > \Delta_{\text{S}}$  PPL)

Metagreywacke  $\text{Nd}_{\text{I}} = 2429$ – $2690 \text{ Ma}$  biotite monzogranite ( $\text{Nd}_{\text{I}} = 3470 \text{ Ma}$  3f).  $\text{Nd}_{\text{I}} = 2095$ – $2310 \text{ Ma}$ , greywacke  $\Delta_{\text{C}} > \Delta_{\text{S}}$  Lake Harbour Group  $\Delta_{\text{L}} = 2500 \text{ Ma}$ ,  $\text{P}_{\text{r}} = 5 \text{ kbar}$ ,  $\text{T}_{\text{p}} = 600 \text{–} 800 \text{ °C}$ . Longstaff Bluff  $\text{Nd}_{\text{I}} = 2500 \text{ Ma}$  Piling Group  $\Delta_{\text{L}} = 2450 \text{ Ma}$  (Wodicka et al., 2014),  $\text{Nd}_{\text{I}} = 2500 \text{ Ma}$  granitoid- $\Delta_{\text{L}} = 2450 \text{ Ma}$ .

**Paleoproterozoic mafic-ultramafic sills ( $\Delta_{\text{C}} > \Delta_{\text{S}}$  PLHu–PLHd)**

$\Delta_{\text{L}} = 2500 \text{ Ma}$  Ld $\sigma^{\text{m}}$   $\Delta_{\text{L}} = 2500 \text{ Ma}$   $\text{Nd}_{\text{I}} = 2429$ – $2690 \text{ Ma}$   $\text{Nd}_{\text{I}} = 2500 \text{ Ma}$   $\text{Nd}_{\text{I}} = 2500 \text{ Ma}$   $\text{Nd}_{\text{I}} = 2500 \text{ Ma}$  ultramafic rocks  $\Delta_{\text{L}} = 2500 \text{ Ma}$   $\Delta_{\text{C}} > \Delta_{\text{S}}$  (Liikane et al., 2015)  $\Delta_{\text{L}} = 2500 \text{ Ma}$  siliciclastic strata  $\Delta_{\text{L}} = 2500 \text{ Ma}$   $\text{Nd}_{\text{I}} = 2500 \text{ Ma}$  mafic  $\Delta_{\text{L}} = 2500 \text{ Ma}$  ultramafic rocks  $\Delta_{\text{L}} = 2500 \text{ Ma}$  their host metasedimentary units,  $\Delta_{\text{L}} = 2500 \text{ Ma}$   $\text{Nd}_{\text{I}} = 2500 \text{ Ma}$  clinopyroxene-orthopyroxene±hornblende metapyroxenite ( $\Delta_{\text{L}} = 2500 \text{ Ma}$  4a),  $\Delta_{\text{L}} = 2500 \text{ Ma}$  olivine-clinopyroxene-orthopyroxene metaperidotite  $\Delta_{\text{L}} = 2500 \text{ Ma}$  metaleucogabbro up section ( $\Delta_{\text{L}} = 2500 \text{ Ma}$  4b).  $\Delta_{\text{L}} = 2500 \text{ Ma}$ ,  $\Delta_{\text{C}} > \Delta_{\text{S}}$  MSc thesis  $\Delta_{\text{L}} = 2500 \text{ Ma}$   $\Delta_{\text{L}} = 2500 \text{ Ma}$  petrology, geochemistry,  $\Delta_{\text{L}} = 2500 \text{ Ma}$  geochronology  $\Delta_{\text{L}} = 2500 \text{ Ma}$   $\text{Nd}_{\text{I}} = 2500 \text{ Ma}$  Raglan Ni-Cu deposit ( $\Delta_{\text{L}} = 2500 \text{ Ma}$  2).

Liikane et al. (2015).

**Paleoproterozoic metaplutonic suite ( $\Delta_{\text{C}} > \Delta_{\text{S}}$  Pg–Psb)**

Gabbro ( $\Delta_{\text{C}} > \Delta_{\text{S}}$  Pg)

$\Delta_{\text{C}} > \Delta_{\text{S}}$  pegmatitic, kilometre-scale, layered biotite-clinopyroxene-magnetite±hornblende gabbro plutons  $\Delta_{\text{C}} > \Delta_{\text{S}}$  Sylvia Grinnell Lake–Clearwater Fiord  $\Delta_{\text{C}} > \Delta_{\text{S}}$ .  $\Delta_{\text{C}} > \Delta_{\text{S}}$  plutons  $\Delta_{\text{C}} > \Delta_{\text{S}}$  clinopyroxene-bearing anorthosite.

Quartz diorite ( $\Delta_{\text{C}} > \Delta_{\text{S}}$  Pg)

Quartz diorite  $\Delta_{\text{C}} > \Delta_{\text{S}}$  biotite-clinopyroxene-orthopyroxene±hornblende.  $\Delta_{\text{C}} > \Delta_{\text{S}}$  plutons  $\Delta_{\text{C}} > \Delta_{\text{S}}$  ( $\Delta_{\text{C}} > \Delta_{\text{S}}$  5a).





assemblage in felsic granitoid units (Weller et al., 2015). CL<sup>o</sup>d<sup>4</sup> ፩<sub>2</sub>OΔ<sup>5</sup><sub>3</sub>SiO<sub>4</sub>CaO<sub>2</sub>Al<sub>2</sub>O<sub>3</sub>MgO<sub>2</sub>FeO<sub>2</sub>Fe<sub>2</sub>O<sub>3</sub> MnO-Na<sub>2</sub>O-CaO-K<sub>2</sub>O-FeO-MgO-Al<sub>2</sub>O<sub>3</sub>-SiO<sub>2</sub>-H<sub>2</sub>O-TiO<sub>2</sub> phase-diagram (Figure 6c) was used to calculate P-T conditions for the samples. The dataset tc-ds55 (Holland and Powell, 1998; Palin et al., 2015) was used to calculate P-T conditions. The calculated P-T conditions are shown in Figure 6d. The calculated P-T conditions are shown in Figure 6d. The calculated P-T conditions are shown in Figure 6d.

A P-T diagram (Figure 6d) shows that the sample (sample 95-D078B from Thériault et al., 2001; Figure 6a) has a monzogranite composition (sample 95-D078B from Thériault et al., 2001; Figure 6a) and high variance assemblages (Figure 6e) in the P-T space, ΔP = 400–800 MPa, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6f) orthopyroxene (Figure 6g) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6h) orthopyroxene (Figure 6i) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6j) orthopyroxene (Figure 6k) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6l) orthopyroxene (Figure 6m) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6n) orthopyroxene (Figure 6o) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6p) orthopyroxene (Figure 6q) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6r) orthopyroxene (Figure 6s) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6t) orthopyroxene (Figure 6u) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6v) orthopyroxene (Figure 6w) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6x) orthopyroxene (Figure 6y) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6z) orthopyroxene (Figure 6aa) sub-solidus conditions.

The P-T diagram (Figure 6d) shows that the sample (sample 95-D078B from Thériault et al., 2001; Figure 6a) has a monzogranite composition (sample 95-D078B from Thériault et al., 2001; Figure 6a) and high variance assemblages (Figure 6e) in the P-T space, ΔP = 400–800 MPa, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6f) orthopyroxene (Figure 6g) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6h) orthopyroxene (Figure 6i) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6j) orthopyroxene (Figure 6k) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6l) orthopyroxene (Figure 6m) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6n) orthopyroxene (Figure 6o) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6p) orthopyroxene (Figure 6q) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6r) orthopyroxene (Figure 6s) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6t) orthopyroxene (Figure 6u) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6v) orthopyroxene (Figure 6w) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6x) orthopyroxene (Figure 6aa) sub-solidus conditions. The P-T range is 400–800°C, ΔT = 700–800°C, metagranitoid rocks, Ldσ<sup>4</sup> magnetite (Figure 6z) orthopyroxene (Figure 6aa) sub-solidus conditions.

orthopyroxene & magnetite (Ca-rich CaMnO<sub>3</sub> - CaFe<sub>2</sub>O<sub>4</sub>) & CaAl<sub>2</sub>O<sub>5</sub> & MnO<sub>2</sub>. The P-T conditions are typical granitoid composition & the pressure varies from 0.0 to 0.5 GPa. The XFe<sup>3+</sup> value is assumed to be 0.1.


The XFe<sup>3+</sup> value varies from 0.0 to 0.5 (Kiss and Tschirhart, 2015a-r). The XFe<sup>3+</sup> value is assumed to be 0.1. The orthopyroxene-bearing metagranitoid has an aeromagnetic signature similar to magnetite-rich plutons (Tschech et al., 2015).

The aeromagnetic signature of orthopyroxene-bearing metagranitoid is similar to magnetite-rich plutons (Tschech et al., 2015). The XFe<sup>3+</sup> value is assumed to be 0.1.

The XFe<sup>3+</sup> value is assumed to be 0.1. The orthopyroxene-bearing metagranitoid has an aeromagnetic signature similar to magnetite-rich plutons (Tschech et al., 2015).







1. **Qallunaat** 1. 0-2015-2002 1967. Geology, Cumberland Sound, District of Franklin; Geological Survey of Canada, Preliminary Map 17-1966, scale 1:506 880. doi:10.4095/108490

1. Blackadar, R.G., 1967. Geology, Cumberland Sound, District of Franklin; Geological Survey of Canada, Preliminary Map 17-1966, scale 1:506 880. doi:10.4095/108490


2. Blackadar, R.G., 1970. Nottingham, Salisbury, and Mill Islands, District of Franklin; Geological Survey of Canada, Map 1205A, scale 1:250 000. doi:10.4095/109212
3. Jackson, G.D., 1971. Operation Penny Highlands, south-central Baffin Island; *in* Report of Activities, Part A: April to October 1970, Geological Survey of Canada, Paper 71-1, Part A, p. 138–140.
4. Trettin, H.P., 1975. Geology, Lower Paleozoic geology, central and eastern parts of Foxe Basin and Baird Peninsula, Baffin Island, District of Franklin; Geological Survey of Canada, Map 1406A, scale 1:500 000. doi:10.4095/109081
5. Morgan, W.C., 1982. Geology, Koch Island, District of Franklin; Geological Survey of Canada, Map 1535A, scale 1:250 000. doi:10.4095/127056
6. Henderson, J.R., 1985a. Geology, McBeth Fiord-Cape Henry Kater, District of Franklin, Northwest Territories; Geological Survey of Canada, Map 1605A, scale 1:250 000. doi:10.4095/120467
7. Henderson, J.R., 1985b. Geology, Ekalugad Fiord-Home Bay, District of Franklin, Northwest Territories; Geological Survey of Canada, Map 1606A, scale 1:250 000. doi:10.4095/120468
8. Jackson, G.D., 1998. Geology, Okoa Bay-Padloping Island area, District of Franklin, Northwest Territories; Geological Survey of Canada, Open File 3532, scale 1:250 000. doi:10.4095/209911
9. St-Onge, M.R., Scott, D.J., and Wodicka, N., 1999a. Geology, Frobisher Bay, Nunavut; Geological Survey of Canada, Map 1979A, scale 1:100 000. doi:10.4095/210833
10. St-Onge, M.R., Scott, D.J., and Wodicka, N., 1999b. Geology, Hidden Bay, Nunavut; Geological Survey of Canada, Map 1980A, scale 1:100 000. doi:10.4095/210835
11. St-Onge, M.R., Scott, D.J., and Wodicka, N., 1999c. Geology, McKellar Bay, Nunavut; Geological Survey of Canada, Map 1981A, scale 1:100 000. doi:10.4095/210836
12. St-Onge, M.R., Scott, D.J., and Wodicka, N., 1999d. Geology, Wright Inlet, Nunavut; Geological Survey of Canada, Map 1982A, scale 1:100 000. doi:10.4095/210840
13. St-Onge, M.R., Scott, D.J., and Wodicka, N., 1999e. Geology, Blandford Bay, Nunavut; Geological Survey of Canada, Map 1983A, scale 1:100 000. doi:10.4095/210837
14. St-Onge, M.R., Scott, D.J., and Wodicka, N., 1999f. Geology, Crooks Inlet, Nunavut; Geological Survey of Canada, Map 1984A, 1:100 000 scale. doi:10.4095/210838
15. St-Onge, M.R., Scott, D.J., and Wodicka, N., 1999g. Geology, White Strait, Nunavut; Geological Survey of Canada, Map 1985A, scale 1:100 000. doi:10.4095/210839
16. Sanford, B.V. and Grant, A.C., 2000. Geological framework of the Ordovician system in the southeast Arctic platform, Nunavut; *in* Geology and Paleontology of the southeast Arctic Platform and southern Baffin Island, Nunavut; Geological Survey of Canada, Bulletin 557, p. 13–38.
17. Jackson, G.D., 2002. Geology, Isurtuq River-Nedlukseak Fiord, Nunavut; Geological Survey of Canada, Open File 4259, scale 1:250 000. doi:10.4095/213304
18. St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005a. Geology, Ikpik Bay, Baffin Island, Nunavut; Geological Survey of Canada, Map 2077A, scale 1:100 000. doi:10.4095/221054
19. St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005b. Geology, Flyway Lake, Baffin Island, Nunavut; Geological Survey of Canada, Map 2078A, scale 1:100 000. doi:10.4095/221091

20. St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005c. Geology, Clyde River, Baffin Island, Nunavut; Geological Survey of Canada, Map 2079A, scale 1:100 000. doi:10.4095/221092
21. St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005d. Geology, Piling Bay, Baffin Island, Nunavut; Geological Survey of Canada, Map 2080A, scale 1:100 000. doi:10.4095/221093
22. St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005e. Geology, Straits Bay, Baffin Island, Nunavut; Geological Survey of Canada, Map 2081A, scale 1:100 000. doi:10.4095/221094
23. St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005f. Geology, Dewar Lakes, Baffin Island, Nunavut; Geological Survey of Canada, Map 2082A, scale 1:100 000. doi:10.4095/221095
24. Jackson, G.D., 2006. Geology, Hantzsch River area, Baffin Island, Nunavut; Geological Survey of Canada, Open File 4202, scale 1:250 000. doi:10.4095/221809
25. St-Onge, M.R., Sanborn-Barrie, M., and Young, M.D., 2007a. Geology, Mingo Lake, Baffin Island, Nunavut; Geological Survey of Canada, Open File 5433, scale 1:250 000. doi:10.4095/224163
26. St-Onge, M.R., Sanborn-Barrie, M., and Young, M.D., 2007b. Geology, Foxe Peninsula, Baffin Island, Nunavut; Geological Survey of Canada, Open File 5434, scale 1:250 000. doi:10.4095/224222
27. Sanborn-Barrie, M., Young, M., Whalen, J., and James, D., 2011a. Geology, Ujuktuk Fiord, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 1 (2nd edition, preliminary), scale 1:100 000. doi:10.4095/289237
28. Sanborn-Barrie, M., Young, M., and Whalen, J., 2011b. Geology, Kingnait Fiord, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 2 (2nd edition, preliminary), scale 1:100 000. doi:10.4095/289238
29. Sanborn-Barrie, M., Young, M., Whalen, J., James, D., and St-Onge, M.R., 2011c. Geology, Touak Fiord, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 3 (2nd edition, preliminary), scale 1:100 000. doi:10.4095/289239
30. Sanborn-Barrie, M. and Young, M., 2013a. Geology, Circle Lake, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 5 (preliminary), scale 1:100 000. doi:10.4095/288929
31. Sanborn-Barrie, M., Young, M., Keim, R., and Hamilton, B., 2013. Geology, Sunneshine Fiord, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 6 (preliminary), scale 1:100 000. doi:10.4095/288931
32. Sanborn-Barrie, M. and Young, M., 2013b. Geology, Padle Fiord, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 37 (preliminary), scale 1:100 000. doi:10.4095/292014
33. Sanborn-Barrie, M. and Young, M., 2013c. Geology, Durban Harbour, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 38 (preliminary), scale 1:100 000. doi:10.4095/292015
34. Sanborn-Barrie, M. and Young, M., 2013d. Geology, Qikiqtarjuaq, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 39 (preliminary), scale 1:100 000. doi:10.4095/292016
35. Jackson, G.D. and Sanborn-Barrie, M., 2014. Geology, Pangnirtung Fiord, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 4 (preliminary), scale 1:100 000. doi:10.4095/288928
36. St-Onge, M.R., Rayner, N.M., Steenkamp, H.M., and Gilbert, C., 2015a. Geology, Terra Nivea, Baffin Island, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 215E (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2015-02E, scale 1:100 000. doi:10.4095/296104


37. St-Onge, M.R., Rayner, N.M., Steenkamp, H.M., and Gilbert, C., 2015b. Geology, Pritzler Harbour, Baffin Island, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 216E (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2015-03E, scale 1:100 000. doi:10.4095/296109
38. St-Onge, M.R., Rayner, N.M., Steenkamp, H.M., and Gilbert, C., 2015c. Geology, Grinnell Glacier, Baffin Island, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 217E (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2015-04E, scale 1:100 000. doi:10.4095/296111
39. Steenkamp, H., Gilbert, C., and St-Onge, M.R., 2016a. Geology, Loks Land, Baffin Island, Nunavut, NTS 25-I (part); Geological Survey of Canada, Canadian Geoscience Map 264 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-01, scale 1:100 000. doi:10.4095/297344
40. Steenkamp, H., Gilbert, C., and St-Onge, M.R., 2016b. Geology, Ward Inlet (south), Baffin Island, Nunavut, NTS 25-O (south) and NTS 25-J (part); Geological Survey of Canada, Canadian Geoscience Map 266 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-02, scale 1:100 000. doi:10.4095/297349
41. Steenkamp, H., Gilbert, C. and St-Onge, M.R., 2016c. Geology, Ward Inlet (north), Baffin Island, Nunavut, NTS 25-O (north); Geological Survey of Canada, Canadian Geoscience Map 265 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-03, scale 1:100 000. doi:10.4095/297348
42. Steenkamp, H., Gilbert, C., and St-Onge, M.R., in press a. Geology, Beekman Peninsula (south), Baffin Island, Nunavut, NTS 25-P (south) and NTS 15-M (part); Geological Survey of Canada, Canadian Geoscience Map 267 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-04, scale 1:100 000. doi:10.4095/297351
43. Steenkamp, H., Gilbert, C., and St-Onge, M.R., in press b. Geology, Beekman Peninsula (north), Baffin Island, Nunavut, NTS 25-P (north) and NTS 15-M (part); Geological Survey of Canada, Canadian Geoscience Map 268 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-05, scale 1:100 000. doi:10.4095/297352
44. Steenkamp, H., Gilbert, C., and St-Onge, M.R., 2016d. Geology, Leybourne Islands (south), Baffin Island, Nunavut, NTS 26-A (south); Geological Survey of Canada, Canadian Geoscience Map 269 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-06, scale 1:100 000. doi:10.4095/297353
45. Steenkamp, H., Gilbert, C., and St-Onge, M.R., 2016e. Geology, Leybourne Islands (north), Baffin Island, Nunavut, NTS 26-A (north); Geological Survey of Canada, Canadian Geoscience Map 271 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-07, scale 1:100 000. doi:10.4095/297355
46. Steenkamp, H., Gilbert, C., and St-Onge, M.R., in press c. Geology, Chidliak Bay (south), Baffin Island, Nunavut, NTS 26-B (south); Geological Survey of Canada, Canadian Geoscience Map 272 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-08, scale 1:100 000. doi:10.4095/297357
47. Steenkamp, H., Gilbert, C., and St-Onge, M.R., in press d. Geology, Chidliak Bay (north), Baffin Island, Nunavut, NTS 26-B (north); Geological Survey of Canada, Canadian Geoscience Map 270 (preliminary); Canada-Nunavut Geoscience Office, Open File Map 2016-09, scale 1:100 000. doi:10.4095/297354








4. ልንግድ ዓሸኑ ቃል ኢትዮጵያ ሌሎች ስልጣን አስተዳደር ሰነዶች ለማዕበት ቅድመ ስለመሳሪያ እንዲያረዳ ተብሎ የመታወቂያ የሚከተሉት ነው፡ የመታወቂያ የሚከተሉት ነው፡ a) የመታወቂያ የሚከተሉት ነው፡ b) የመታወቂያ የሚከተሉት ነው፡ c) የመታወቂያ የሚከተሉት ነው፡ d) የመታወቂያ የሚከተሉት ነው፡



**Figure 5.** ፩. የጂደልናይሮን ማኅበናዎች መግላጭዎች ማኅበናዎች Sylvia Grinnell Lake-Clearwater Fiord-ገር. በስራው እነዚያዎች ተከተል ያለፈ ሆኖ ተመራሽ ስምምነት ይከተል. a) ቨአዲነናይሮን ለሰነድ ለኩል ለተደሏል ማኅበናው ማኅበናዎች ማኅበናዎች የሚመለከውን ዘመንና ሥርዓት. b) የሀሳታዊ ዕቅድ መግላጭ የክ-ፍልspar መግላጭ ማኅበናዎች የሚመለከውን ዘመንና ሥርዓት. c) አላማውን በአዲነናይሮን ለሰነድ ለሚመለከውን ማኅበናዎች የሚመለከውን ዘመንና ሥርዓት. d) የአዲነናይሮን ተከተል ያለፈ ማኅበናዎች ማኅበናዎች የሚመለከውን ዘመንና ሥርዓት. e) የሀሳታዊ ዕቅድ መግላጭ የሚመለከውን ዘመንና ሥርዓት. f) የአዲነናይሮን ተከተል ያለፈ ማኅበናዎች ማኅበናዎች የሚመለከውን ዘመንና ሥርዓት.



6. 95-D078B assemblages (all fields + Grt, Ilm, Kfs, Pl, Qz) at 7 kbar monzogranite, metagranitoid amphibolite-granulite assemblages: a) P-T; b) T- $\text{MH}_2\text{O}$ ; c) T- $X\text{Fe}^{3+}$ ; d) magnetic anomalies biotite±magnetite±orthopyroxene monzogranite.

## ‘b▷zR▷‘A‘b‘c

Blackadar, R.G., 1967. Geological Reconnaissance, southern Baffin Island, District of Franklin; Geological Survey of Canada, Paper 66-47, 32 p. doi:10.4095/100926

Buchan, K.L. and Ernst, R.E., 2013. Diabase dyke swarms of Nunavut, Northwest Territories and Yukon, Canada; Geological Survey of Canada, Open File 7464. doi:10.4095/293149

Chadwick, T.C., St-Onge, M.R., Weller, O.M., Carr, S.D., and Dyck, B.J., 2015. Ptarmigan Fiord basement-cover thrust imbricates, Baffin Island, Nunavut: summary of fieldwork; *in* Summary of Activities 2015, Canada-Nunavut Geoscience Office, p. 61–72.

Clarke, D.B., Cameron, B.I., Muecke, G.K., and Bates, J.L., 1989. Early Tertiary basalts from the Labrador Sea and Davis Strait region; Canadian Journal of Earth Sciences, v. 26, p. 956–968.

Corrigan, D., Pehrsson, S., Wodicka, N., and de Kemp, E., 2009. The Palaeoproterozoic Trans-Hudson Orogen: a prototype of modern accretionary processes; *in* Ancient Orogenes and Modern Analogues, J.B. Murphy, J.D. Keppie, and A.J. Hynes (ed.), The Geological Society, London, Special Publications, v. 327, p. 457–479. doi:10.1144/SP327.19

Dunphy, J.M. and Ludden, J.N., 1998. Petrological and geochemical characteristics of a Paleoproterozoic magmatic arc (Narsajuaq Terrane, Ungava Orogen, Canada) and comparisons to Superior Province granitoids; Precambrian Research, v. 91, p. 109–142.

Dyck, B.J. and St-Onge, M.R., 2014. Dehydration-melting reactions, leucogranite emplacement and the Paleoproterozoic structural evolution of Hall Peninsula, Baffin Island, Nunavut; *in* Summary of Activities 2013, Canada-Nunavut Geoscience Office, p. 73–84.

Heaman, L.M., LeCheminant, A.N., and Rainbird, R.H., 1992. Nature and timing of Franklin igneous events, Canada: implications for a Late Proterozoic mantle plume and the break-up of Laurentia; Earth and Planetary Science Letters, v. 109, p. 117–131.

Henderson, J.R., 1985. Geology, McBeth Fiord-Cape Henry Kater, District of Franklin, Northwest Territories; Geological Survey of Canada, Map 1605A, scale 1:250 000. doi:10.4095/120467

Hoffman, P.F., 1988. United Plates of America, the birth of a craton: Early Proterozoic assembly and growth of Laurentia; Annual Reviews of Earth and Planetary Sciences, v. 16, p. 543–603.

Holland, T.J.B. and Powell, R., 1998. An internally consistent thermodynamic dataset for phases of petrological interest; Journal of Metamorphic Geology, v. 16, p. 309–343.

Kiss, F. and Tschirhart, V., 2015a. Residual total magnetic field, aeromagnetic survey of the Amittoq Lake area, Baffin Island, Nunavut, NTS 26-J south; Geological Survey of Canada, Open File 7888, scale 1:100 000. doi:10.4095/296504

Kiss, F. and Tschirhart, V., 2015b. First vertical derivative of the magnetic field, aeromagnetic survey of the Amitto Lake area, Baffin Island, Nunavut, NTS 26-J south; Geological Survey of Canada, Open File 7889, scale 1:100 000. doi:10.4095/296505

Kiss, F. and Tschirhart, V., 2015c. Residual total magnetic field, aeromagnetic survey of the Amitto Lake area, Baffin Island, Nunavut, NTS 26-K south; Geological Survey of Canada, Open File 7890, scale 1:100 000. doi:10.4095/296506

Kiss, F. and Tschirhart, V., 2015d. First vertical derivative of the magnetic field, aeromagnetic survey of the Amitto Lake area, Baffin Island, Nunavut, NTS 26-K south; Geological Survey of Canada, Open File 7891, scale 1:100 000. doi:10.4095/296507

Kiss, F. and Tschirhart, V., 2015e. Residual total magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-B/North, Nunavut; Geological Survey of Canada, Open File 7819, scale 1:100 000. doi:10.4095/296382

Kiss, F. and Tschirhart, V., 2015f. First vertical derivative of the magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-B/North, Nunavut; Geological Survey of Canada, Open File 7820, scale 1:100 000. doi:10.4095/296383

Kiss, F. and Tschirhart, V., 2015g. Residual total magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-C/South, Nunavut; Geological Survey of Canada, Open File 7821, scale 1:100 000. doi:10.4095/296384

Kiss, F. and Tschirhart, V., 2015h. First vertical derivative of the magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-C/South, Nunavut; Geological Survey of Canada, Open File 7822, scale 1:100 000. doi:10.4095/296385

Kiss, F. and Tschirhart, V., 2015i. Residual total magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-C/North, Nunavut; Geological Survey of Canada, Open File 7823, scale 1:100 000. doi:10.4095/296386

Kiss, F. and Tschirhart, V., 2015j. First vertical derivative of the magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-C/North, Nunavut; Geological Survey of Canada, Open File 7824, scale 1:100 000. doi:10.4095/296387

Kiss, F. and Tschirhart, V., 2015k. Residual total magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-F/South, Nunavut; Geological Survey of Canada, Open File 7825, scale 1:100 000. doi:10.4095/296388

Kiss, F. and Tschirhart, V., 2015l. First vertical derivative of the magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-F/South, Nunavut; Geological Survey of Canada, Open File 7826, scale 1:100 000. doi:10.4095/296389

Kiss, F. and Tschirhart, V., 2015m. Residual total magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-F/North, Nunavut; Geological Survey of Canada, Open File 7827, scale 1:100 000. doi:10.4095/296390

Kiss, F. and Tschirhart, V., 2015n. First vertical derivative of the magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-F/North, Nunavut; Geological Survey of Canada, Open File 7828, scale 1:100 000. doi:10.4095/296391

Kiss, F. and Tschirhart, V., 2015o. Residual total magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-G/South, Nunavut; Geological Survey of Canada, Open File 7829, scale 1:100 000. doi:10.4095/296392

Kiss, F. and Tschirhart, V., 2015p. First vertical derivative of the magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-G/South, Nunavut; Geological Survey of Canada, Open File 7830, scale 1:100 000. doi:10.4095/296393

Kiss, F. and Tschirhart, V., 2015q. Residual total magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-G/North, Nunavut; Geological Survey of Canada, Open File 7831, scale 1:100 000. doi:10.4095/296394

Kiss, F. and Tschirhart, V., 2015r. First vertical derivative of the magnetic field, aeromagnetic survey of the McKeand River area, part of NTS 26-G/North, Nunavut; Geological Survey of Canada, Open File 7832, scale 1:100 000. doi:10.4095/296395

Lesher, C.M., 2007. Ni-Cu-(PGE) deposits in the Raglan area, Cape Smith Belt, New Quebec; *in* Mineral Deposits of Canada: a Synthesis of Major Deposit Types, District Metallogeny, the Evolution of Geological Provinces and Exploration Methods, W.D. Goodfellow (ed.), Geological Association of Canada, Special Publication, v. 5, p. 351–386.

Liikane, D.A., St-Onge, M.R., Kjarsgaard, B.A., Rayner, N.M., Ernst, R.E., and Kastek, N., 2015. Frobisher suite mafic, ultramafic and layered mafic-ultramafic sills, southern Baffin Island; *in* Summary of Activities 2015, Canada-Nunavut Geoscience Office, p. 21–32.

Lucas, S.B. and Byrne, T., 1992. Footwall involvement during arc-continent collision, Ungava orogen, northern Canada; Journal of the Geological Society of London, v. 149, p. 237–248.

Lucas, S.B. and St-Onge, M.R., 1992. Terrane accretion in the internal zone of the Ungava orogen, northern Quebec. Part 2: Structural and metamorphic history; Canadian Journal of Earth Sciences, v. 29, p. 765–782.

Nekvasil, H., 1991. Ascent of felsic magmas and formation of rapakivi; American Mineralogist, v. 76, p. 1279–1290.

Palin, RM., Weller, O.M., Waters, D.J., and Dyck, B., 2015. Quantifying geological uncertainty in metamorphic phase equilibria modelling; a Monte Carlo assessment and implications for tectonic interpretations; Geoscience Frontiers. doi:10.1016/j.gsf.2015.08.005

Powell, R. and Holland, T.J.B., 2008. On thermobarometry; Journal of Metamorphic Geology, v. 26, p. 155–179.

Scott, D.J., 1997. Geology, U-Pb, and Pb-Pb geochronology of the Lake Harbour area, southern Baffin Island: implications for the Paleoproterozoic tectonic evolution of north-eastern Laurentia; Canadian Journal of Earth Sciences, v. 34, p. 140–155.

Scott, D.J. and Wodicka, N., 1998. A second report on the U-Pb geochronology of southern Baffin Island; Geological Survey of Canada, Current Research 1998-F, p. 47–57.

Scott, D.J., St-Onge, M.R., Wodicka, N., and Hanmer, S., 1997. Geology of the Markham Bay – Crooks Inlet area, southern Baffin Island, Northwest Territories; Geological Survey of Canada, Current Research 1997-C, p. 157–166.

St-Onge, M.R. and Lucas, S.B., 1994. Controls on the regional distribution of iron-nickel-copper-platinum group element sulfide mineralization in the eastern Cape Smith Belt, Quebec; Canadian Journal of Earth Sciences, v. 31, p. 206–218.

St-Onge, M.R., Hanmer, S., and Scott, D.J., 1996. Geology of the Meta Incognita Peninsula, south Baffin Island: tectonostratigraphic units and regional correlations; Geological Survey of Canada, Current Research 1996-C, p. 63–72.

St-Onge, M.R., Scott, D.J. and Lucas, S.B., 2000a. Early partitioning of Quebec: Microcontinent formation in the Paleoproterozoic; Geology, v. 28, p. 323–326.

St-Onge, M.R., Scott, D.J., Wodicka, N., and Lucas, S.B., 1998. Geology of the McKellar Bay – Wight Inlet – Frobisher Bay area, southern Baffin Island, Northwest Territories; Geological Survey of Canada, Current Research 1998-C, p. 43–53.

St-Onge, M.R., Searle, M.P., and Wodicka, N., 2006. Trans-Hudson Orogen of North America and Himalaya-Karakoram-Tibetan Orogen of Asia: Structural and thermal characteristics of the lower and upper plates; Tectonics, v. 25, TC4006, 22p. doi:10.1029/2005TC001907

St-Onge, M.R., Wodicka, N., and Ijewliw, O., 2007. Polymetamorphic evolution of the Trans-Hudson Orogen, Baffin Island, Canada: Integration of petrological, structural and geochronological data; Journal of Petrology, v. 48, p. 271–302. doi:10.1093/petrology/eg1060

St-Onge, M.R., Wodicka, N., and Lucas, S.B., 2000b. Granulite- and amphibolite-facies metamorphism in a convergent plate-margin setting: Synthesis of the Quebec-Baffin segment of Trans-Hudson Orogen. Canadian Mineralogist, v. 38, p. 379–398.

St-Onge, M.R., Van Gool, J.A.M., Garde, A.A., and Scott, D.J., 2009. Correlation of Archaean and Palaeoproterozoic units between northeastern Canada and western Greenland: constraining the pre-collisional upper plate accretionary history of the Trans-Hudson orogen; *in* Earth Accretionary Systems in Space and Time, P.A. Cawood and A. Kröner, The Geological Society, London, Special Publications, v. 318, p. 193–235. doi:10.1144/SP318.7

Thériault, R.J., St-Onge, M.R., and Scott, D.J., 2001. Nd isotopic and geochemical signature of the Paleoproterozoic Trans-Hudson Orogen, southern Baffin Island, Canada: implications for the evolution of eastern Laurentia; *Precambrian Research*, v. 108, p. 113–138.

Tschirhart, V., St-Onge, M.R., and Weller, O.M., 2015. Preliminary geophysical interpretation of the McKeand River area, Baffin Island, Nunavut: insights from gravity, magnetic and geological data; *in* Summary of Activities 2015, Canada-Nunavut Geoscience Office, p. 49–60.

Weller, O.M., Dyck, B.J., St-Onge, M.R., Rayner, N.M., and Tschirhart, V., 2015. Completing the bedrock mapping of southern Baffin Island, Nunavut: plutonic suites and regional stratigraphy; *in* Summary of Activities 2015, Canada-Nunavut Geoscience Office, p. 33–48.

Weller, O.M., St-Onge, M.R., Rayner, N., Searle, M.P., and Waters, D.J., 2016. Miocene magmatism in the Western Nyainqntanglha mountains of southern Tibet: an exhumed bright spot?; *Lithos*, doi:10.1016/j.lithos.2015.06.024

Weller, O.M., St-Onge, M.R., Searle, M.P., Rayner, N., Waters, D.J., Chung, S.L., Palin, R.M., Lee, Y.H., and Xu, X.W., 2013. Quantifying Barrovian metamorphism in the Danba Structural Culmination of eastern Tibet; *Journal of Metamorphic Geology*, v. 31, p. 909–935. doi: 10.1111/jmg.12050

Whalen, J.B., Wodicka, N., Taylor, B.E., and Jackson, G.D., 2010. Cumberland batholith, Trans-Hudson Orogen, Canada: Petrogenesis and implications for Paleoproterozoic crustal and orogenic processes; *Lithos*, v. 117, p. 99–118. doi:10.1016/j.lithos.2010.02.008

Wodicka, N. and Scott, D.J., 1997. A preliminary report on the U-Pb geochronology of the Meta Incognita Peninsula, southern Baffin Island, Northwest Territories; *Geological Survey of Canada, Current Research* 1997-C, p. 167–178.

Wodicka, N., St-Onge, M.R., Corrigan, D., Scott, D.J., and Whalen, J.B., 2014. Did a proto-ocean basin form along the southeastern Rae cratonic margin? Evidence from U-Pb geochronology, geochemistry (Sm-Nd and whole rock), and stratigraphy of the Paleoproterozoic Piling Group, northern Canada; *GSA Bulletin*, v. 126, p. 1625–1653. doi: 10.1130/B31028.1



ԿԱՐԼԾԱ ԽՄԼ ԴԵՅՎԱԼԳԼ ՀԿԱՐԱՎԾ ԼՇԱԾԵՐԿԿ ՀՃԵԾԱ  
ՆԵՄԱԾԱԾՐԱԾ. ՀԿԱԾԵՐՎԱԾԱ, ԵՄԱԾՎԾԱ ՈՈԴԿԱԼԾ ԱՆԴԵԿԿ  
ՆԺԱՒԺԱԾԿԾ ԿԲԿՐԱՔՈՐԱԾ ԵՄԱԾՎԾԱ ՐՈԼԸ.

## **ՀԱԿ/ՀԿԱ ՀԵՐԱՎՐԱԾ ԵՎ ԿԺՈՎԾ ՏԵՂՅԱԾԸ ՀԱՅ ԻՇՈՎԾԸ.**

### **LICENCE AGREEMENT**

View the licence agreement at

<http://open.canada.ca/en/open-government-licence-canada>

### **ACCORD DE LICENCE**

Voir l'accord de licence à

<http://ouvert.canada.ca/fr/licence-du-gouvernement-ouvert-canada>