



Natural Resources
Canada

Ressources naturelles
Canada

**GEOLOGICAL SURVEY OF CANADA
OPEN FILE 8206**

**Combined depth and S1 maps from Rock-Eval 6/TOC data
of the Canol Formation, northern Mackenzie Valley,
Northwest Territories**

A.M. Durbano, T. Hadlari, K.M. Fallas, and C. Jiang

2017



Canada 



GEOLOGICAL SURVEY OF CANADA OPEN FILE 8206

Combined depth and S1 maps from Rock-Eval 6/TOC data of the Canol Formation, northern Mackenzie Valley, Northwest Territories

A.M. Durbano, T. Hadlari, K.M. Fallas, and C. Jiang

2017

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources, 2017

Information contained in this publication or product may be reproduced, in part or in whole, and by any means, for personal or public non-commercial purposes, without charge or further permission, unless otherwise specified.

You are asked to:

- exercise due diligence in ensuring the accuracy of the materials reproduced;
- indicate the complete title of the materials reproduced, and the name of the author organization; and
- indicate that the reproduction is a copy of an official work that is published by Natural Resources Canada (NRCan) and that the reproduction has not been produced in affiliation with, or with the endorsement of, NRCan.

Commercial reproduction and distribution is prohibited except with written permission from NRCan. For more information, contact NRCan at nrcan.copyrightdroitdauteur.nrcan@canada.ca.

doi:10.4095/299885

This publication is available for free download through GEOSCAN (<http://geoscan.nrcan.gc.ca/>).

Recommended citation

Durbano, A.M., Hadlari, T., Fallas K.M., and Jiang, C., 2017. Combined depth and S1 maps from Rock-Eval 6/TOC data of the Canol Formation, northern Mackenzie Valley, Northwest Territories; Geological Survey of Canada, Open File 8206, 1 .zip file. doi:10.4095/299885

ABSTRACT

Previously published Rock-Eval 6/TOC pyrolysis data are used as a proxy to map the geographic distribution of hydrocarbons in the Canol Formation. Mainly from wells, S1 is interpolated with Kriging software. The contoured representation of both S1 and S1/TOC are overlain by a depth map of the Canol Formation, which was generated from interpretations of seismic data and well logs and is contained within a subsurface zero edge determined from both subsurface data and bedrock maps. The Norman Wells area shows relatively high S1 and S1/TOC values and areas structurally along strike also contain relatively high S1/TOC values.

INTRODUCTION

Rock-Eval 6/TOC pyrolysis data is a fairly standard technique for measuring source rock potential and S1 values in particular represent free hydrocarbons in the rock that either were generated from kerogen in the source rock itself since deposition or accumulated by migration. They are important indicators for the prospectivity of shale rocks as unconventional hydrocarbon plays. When considering unconventional shale oil plays, the presence of natural fractures can have a negative influence on oil production from stimulated hydraulic fractures (Gale et al., 2007). Pre-existing fractures will ultimately focus hydraulic fluids and decrease the effectiveness of hydraulic fracture treatments. In the case of the Canol Formation, its lithology is unique in that it is very siliceous (see Pyle et al., 2011) compared to other Devonian mudstones in the region and could therefore behave in a more brittle manner than typical mudstone. Though no rheological analysis of the Canol Formation was performed we note the empirical observation that in the Norman Wells area that the Canol Formation contains natural fractures at reservoir depths of 500-350 m (Yose et al., 2001; see discussion in Hadlari, 2015). It stands to reason that the Canol Formation has better prospectivity for unconventional hydrocarbons at greater depths and the deeper it is the more likely that stimulated fractures will be oriented vertically (Hubbert and Willis, 1972). This report uses previously published Rock-Eval 6/TOC pyrolysis data of the Canol Formation as a proxy to interpolate a surface representing the geographic distribution of hydrocarbons in this unconventional shale oil play with superimposed depth contours to identify areas of prospectivity.

METHODOLOGY

The source of Rock-Eval 6/TOC pyrolysis data shown here for the Canol Formation is a compilation by Pyle et al. (2015). In the case of multiple samples for a given outcrop or well location, the results were averaged. Spatial S1 data were kriged in ArcGIS using the Geostatistical Analyst plug-in. Canol Formation depths were derived from an interpretation of seismic data and exploration wells that was developed in the manner described in MacLean (2012). The distribution of the Canol Formation in the subsurface is approximated by the distribution of the entire Horn River Group, of which the Canol Formation is the upper stratigraphic unit, and integrates both subsurface and bedrock maps (Aitken et al., 1969; Aitken and Cook, 1979a, b; Cook and Aitken, 1970, 1975a, 1975b, 1975c; Fallas, 2013; Fallas and MacLean, 2013; Fallas and MacNaughton 2013, 2014a, b; Fallas et al., 2013a,b,c; Norris, 1981a, b, c, d, 1982). Surface fault data is sourced from a 1:5 000 000 compilation by Wheeler et al. (1996) and 1:50 000 to 1:100 000 maps by Fallas (2013a, b, c, d), Fallas and MacLean (2013a, b), Fallas and MacNaughton (2013, 2014a, b, c) and Fallas et al. (2013a, b, c, d). Subsurface faults are sourced from MacLean (2012).

RESULTS

The contoured depth map of the Canol Formation with interpolated S1 contours in Figure 1 shows the geographic distribution of free hydrocarbons indicated by Rock-Eval pyrolysis. In general the central Mackenzie area has relatively high S1 values whereas free hydrocarbon contents decrease toward the northern and southern edges of the study area. Some of the highest S1 values are recorded near Norman Wells, consistent with early reports of oil production from the Canol Formation (see Irish and Kempthorne, 1987).

By normalizing the S1 values to bulk TOC, areas with anomalously high S1 are indicated by the S1/TOC map (Figure 2). The interpolated S1/TOC contours in Figure 2 are superimposed with a contoured depth map of the Canol Formation. In the central part of the study area where S1 is elevated, the S1/TOC map further delimits smaller regions with anomalously high S1 relative to TOC. Relatively high S1/TOC values at Norman Wells indicate that this method has potential to identify a Norman Wells-type play. It is tempting to suggest that the geographic trend of high S1/TOC follows along the strike of faults, in the region of elevated S1, but more work is required to explore this relationship.

It is intended that maps of free hydrocarbons in Canol Formation rocks as measured by Rock-Eval pyrolysis inform assessment of the unconventional resource potential in the Central Mackenzie region.

ACKNOWLEDGEMENTS

This work benefited from discussions with Dr. Martin Fowler. It follows extensive subsurface work in the region by Bernie MacLean. The manuscript was reviewed by Dr. Zhuoheng Chen.

REFERENCES

- Aitken, J.D., Ayling, M.E., Balkwill, H.R., Cook, D.G., Mackenzie, W.S., and Yorath, C.J., 1969. Fort Good Hope, District of Mackenzie; Geological Survey of Canada, Preliminary Map 4-1969; 1 sheet, doi:10.4095/109213.
- Aitken, J.D. and Cook, D.G., 1979a. Geology, Upper Ramparts River, District of Mackenzie; Geological Survey of Canada, Map 1452A; 1 sheet, doi:10.4095/123318.
- Aitken, J.D. and Cook, D.G., 1979b. Geology, Sans Sault Rapids, District of Mackenzie; Geological Survey of Canada, Map 1453A; 1 sheet, doi:10.4095/123316.
- Cook, D.G. and Aitken, J.D., 1970. Geology, Colville Lake and Coppermine, District of Mackenzie; Geological Survey of Canada, Preliminary Map 12-1970; 1 sheet, doi:10.4095/220386.
- Cook, D.G. and Aitken, J.D., 1975a. Geology, Ontaratue River, District of Mackenzie; Geological Survey of Canada, Map 1408A; 1 sheet, doi:10.4095/109027.
- Cook, D.G. and Aitken, J.D., 1975b. Geology, Travaillant Lake, District of Mackenzie; Geological Survey of Canada, Map 1409A; 1 sheet, doi:10.4095/109028.
- Cook, D.G. and Aitken, J.D., 1975c. Geology, Canot Lake, District of Mackenzie; Geological Survey of Canada, Map 1410A; 1 sheet, doi:10.4095/109029.
- Cook, D.G. and MacLean, B.C., 1999. The Imperial anticline, a fault-bend fold above a bedding-parallel thrust ramp, Northwest Territories, Canada. *Journal of Structural Geology*, 21, p. 215-228.
- Fallas, K.M., 2013a. Geology, Mahony Lake (northwest), Northwest Territories; Geological Survey of Canada, Canadian Geoscience Map 88, scale 1:100 000. doi:10.4095/292211.
- Fallas, K.M., 2013b. Geology, Mahony Lake (southeast), Northwest Territories; Geological Survey of Canada, Canadian Geoscience Map 90, scale 1:100 000. doi:10.4095/292282.
- Fallas, K.M., 2013c. Geology, Norman Wells (northeast), Northwest Territories; Geological Survey of Canada, Canadian Geoscience Map 99, scale 1:100 000. doi:10.4095/292291.
- Fallas, K.M., 2013d. Geology, Norman Wells (northwest), Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 98, 2013; 1 sheet, doi:10.4095/292290.
- Fallas, K.M. and MacLean, B.C., 2013a. Geology, Fort Norman (southwest), Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 93, 2013; 1 sheet, doi:10.4095/292285.
- Fallas, K.M. and MacLean, B.C., 2013b. Geology, Mahony Lake (northeast), Northwest Territories; Geological Survey of Canada, Canadian Geoscience Map 89, scale 1:100 000. doi:10.4095/292174.
- Fallas, K.M. and MacNaughton, R B., 2013. Geology, Norman Wells (southeast), Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 100, 2013; 1 sheet, doi:10.4095/292292.

Fallas, K.M. and MacNaughton, R.B., 2014a. Geology, Carcajou Canyon (northwest), Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 94, 2014; 1 sheet, doi:10.4095/292286.

Fallas, K.M. and MacNaughton, R.B., 2014b. Geology, Carcajou Canyon (southeast), Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 96, 2014; 1 sheet, doi:10.4095/292288.

Fallas, K.M. and MacNaughton, R.B., 2014c. Geology, Carcajou Canyon (southwest), Northwest Territories; Geological Survey of Canada, Canadian Geoscience Map 97, scale 1:100 000. doi:10.4095/292289

Fallas, K.M., Hadlari, T. and MacLean, B.C., 2013a. Geology, Carcajou Canyon (northeast), Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 95, 2013; 1 sheet, doi:10.4095/292287.

Fallas, K.M., MacLean, B.C., and Hadlari, T., 2013b. Geology, Mahony Lake (southwest), Northwest Territories; Geological Survey of Canada, Canadian Geoscience Map 91, scale 1:100 000. doi:10.4095/292283

Fallas, K.M., MacLean, B.C., and Proks, T., 2013c. Geology, Fort Norman (northwest), Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 92, 2013; 1 sheet, doi:10.4095/292284.

Fallas, K.M., MacNaughton, R.B., MacLean, B.C., and Hadlari, T., 2013d. Geology, Norman Wells (southwest), Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 101, 2013; 1 sheet, doi:10.4095/292293.

Feinstein, S., Brooks, P.W., Gentzis, T., Goodarzi, F., Snowdon, L.R., and Williams, G.K., 1988b. Thermal maturity in the Mackenzie Corridor, Northwest and Yukon Territories; Canada; Geological Survey of Canada, Open File 1944.

Fowler, M.G., Obermajer M., and Stasiuk, L.D., 2002. Rock-Eval/TOC data for Devonian potential source rocks, Western Canada Sedimentary Basin. Geological Survey of Canada, Open File 1579, 1 CDROM.

Gal, L.P., Allen, T.L., Hadlari, T., and Zantvoort, W.G. Chapter 10 – Petroleum System Elements. In: Regional Geoscience Studies and Petroleum Potential, Peel Plateau and Plain: Project Volume, edited by L.J. Pyle and A.L. Jones, Northwest Territories Geoscience Office, NWT Open File 2009-02, p. 477-549.

Gale, J.F.W., Reed, R.M., and Holder, J., 2007. Natural fractures in the Barnett Shale and their importance for hydraulic fracture treatments. AAPG Bulletin, v. 91, p. 603-622.

Grass, D.B., 1988. Surface geological field program, lower Mackenzie Valley, Northwest Territories. Report on geological examination of Devonian outcrop, Mackenzie River Valley and Mackenzie Mountains, NWT; Amoco Canada Petroleum Company Limited, National Energy Board Report #9233-A4-2E.

Hadlari, T. 2015. Oil migration driven by exhumation of the Canol Formation oil shale: a new conceptual model for the Norman Wells oil field, northwestern Canada. *Marine and Petroleum Geology* 65, p. 172-177.

Hubbert, M.K. and Willis, D.G., 1972. Mechanics of hydraulic fracturing. *AAPG, Memoir* 18, p. 239-257.

Irish, J.P.R., Kempthorne, R.H., 1987. The study of natural fractures in a reef complex, Norman Wells oil field, Canada. In: *Reservoir Sedimentology*, edited by R.W. Tillman, K.J. Weber, Society of Economic Paleontologists and Mineralogists, Special Publication 40, pp. 153-168.

Lemieux, Y., MacLean, B.C., and Pyle, L.J., 2009. Chapter 2 – Regional structure of Peel Plateau and Plain. In: *Regional Geoscience Studies and Petroleum Potential, Peel Plateau and Plain: Project Volume*, edited by L.J. Pyle and A.L. Jones, Northwest Territories Geoscience Office, NWT Open File 2009-02, p. 43-82.

MacLean, B.C., 2012. GIS-enabled structure maps of subsurface Phanerozoic strata, northwestern Northwest Territories. Geological Survey of Canada, Open File 7172, doi:10.4095/292152.

Norris, D.K., 1981a. Geology, Fort Mcpherson, District of Mackenzie; Geological Survey of Canada, Map 1520A; 1 sheet, doi:10.4095/109651.

Norris, D.K., 1981b. Geology, Arctic Red River, District of Mackenzie; Geological Survey of Canada, Map 1521A; 1 sheet, doi:10.4095/109697.

Norris, D.K., 1981c. Geology, Trail River, Yukon-Northwest Territories; Geological Survey of Canada, Map 1524A; 1 sheet, doi:10.4095/119405.

Norris, D.K., 1981d. Geology, Martin House, Yukon-Northwest Territories; Geological Survey of Canada, Map 1525A; 1 sheet, doi:10.4095/119402.

Norris, D.K., 1982. Geology, Snake River, Yukon-Northwest Territories; Geological Survey of Canada, Map 1592A; 1 sheet, doi:10.4095/109233.

Pyle, L.J., Gal, L.P., and Fiess, K.M., 2014. Devonian Horn River Group: A Reference Section, Lithogeochemical Characterization, Correlation of Measured Sections and Wells, and Petroleum-Potential Data, Mackenzie Plain area (NTS 95M, 95N, 96C, 96D, 96E, 106H, and 106I), NWT; Northwest Territories Geoscience Office, NWT Open File 2014-06, 67 p.

Pyle, L.J., Gal, L.P., and Hadlari, T., 2015. Thermal maturity trends for Horn River Group units and equivalent strata in the Mackenzie Corridor, Northwest Territories and Yukon. Geological Survey of Canada Open File 7850, p. x-x, in press.

Pyle, L.J., Gal, L.P., and Lemiski, R.T., 2011. Measured sections and petroleum potential data (conventional and unconventional) of Horn River Group Outcrops – Part 1, NTS 96D, 96E, and 106H, Northwest Territories. Northwest Territories Geoscience Office, NWT Open File 2011-09, 116 p.

Pyle, L.J., Hadlari, T., Bartolini, M., and Sanei, H., 2014. Measured porosity in Horn River Group outcrop samples, Central Mackenzie Valley, NWT. Northwest Territories Geoscience Office, NWT Open Report 2014-011, 29 p.

Wheeler, J.O., Hoffman, P.F., Card, K.D., Davidson, A., Sandford, B.V., Okulitch, A.V., and Roest, W.R. 1996. Geological map of Canada. Geological Survey of Canada, Map 1860A, scale 1: 5 000 000

Yose, L.A., Brown, S., Davis, T.L., Eiben, T., Kompanik, G.S., and Maxwell, S.R., 2001. 3-D geologic model of a fractured carbonate reservoir, Norman Wells Field, NWT, Canada. Bulletin of Canadian Petroleum Geology 49, p. 86-116.

LIST OF FIGURES

Figure 1. Map showing contoured depth of the Canol Formation, faults and Rock-Eval S1 values. Depths are interpreted from seismic data and well logs (MacLean, 2012). Dark grey faults taken from Wheeler et al. (1996). Black Faults taken from Fallas (2013a, b, c, d), Fallas and MacLean (2013a, b), Fallas and MacNaughton (2013, 2014a, b, c) and Fallas et al. (2013a, b, c, d). Dark blue subsurface faults are interpreted from seismic data (MacLean, 2012). S1 values are from the compilation by Pyle et al. (2015).

Figure 2. Map showing contoured depth of the Canol Formation, faults and Rock-Eval S1/TOC ratios. Depths are interpreted from seismic data and well logs (MacLean, 2012). Dark grey faults taken from Wheeler et al. (1996). Black Faults taken from Fallas (2013a, b, c, d), Fallas and MacLean (2013a, b), Fallas and MacNaughton (2013, 2014a, b, c) and Fallas et al. (2013a, b, c, d). Dark blue subsurface faults are interpreted from seismic data (MacLean, 2012). S1 and TOC values are from the compilation by Pyle et al. (2015).

LIST OF TABLES

Table 1. Tmax values from well and outcrop samples, modified by adding S1/TOC to the compilation by Pyle et al. (2015).