

**References**

Deblonde, C., Cocking, R.B., Kerr, D.E., Campbell, J.E., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Parent, M., Plouffe, A., Robertson, L., Smith, I.R., and Weatherston, A., 2018, Surficial Data Model: the science language of the integrated Geological Survey of Canada data model for surficial geology maps; Geological Survey of Canada, Open File 8236, ver. 2.3.14, 1 .zip file. <https://doi.org/10.4095/308178>

Thomas, R.D. and Dyke, A.S., 1982, Surficial geology, lower Hayes River, District of Keewatin; Geological Survey of Canada, Map 7-1981, scale 1:250 000. <https://doi.org/10.4095/109351>


Thomas, R.D. and Dyke, A.S., 1982, Surficial geology, lower Hayes River, District of Keewatin; Geological Survey of Canada, Map 7-1981, scale 1:250 000. <https://doi.org/10.4095/109351>

**Abstract**

This new surficial geology map product represents the conversion of Preliminary Map 7-1981 (Thomas and Dyke, 1982) and its legend, using the Geological Survey of Canada's Surficial Data Model (SDM version 2.3.14) (Deblonde et al., 2018). All geoscience knowledge and information from Preliminary Map 7-1981 that conformed to the current SDM were maintained during the conversion process. The purpose of converting legacy map data to a common science language and common legend is to enable and facilitate the efficient digital compilation, interpretation, management, and dissemination of geological map information in a structured and consistent manner. This provides an effective knowledge-management tool designed around a geodatabase that can expand, following the type of information to appear on new surficial geology maps.

**Résumé**

Ce nouveau produit cartographique de la géologie des formations superficielles correspond à la conversion de la Carte préliminaire 7-1981 (Thomas et Dyke, 1982) et de sa légende, en se servant du Modèle de données pour les formations superficielles (MDFS version 2.3.14) de la Commission géologique du Canada (Deblonde et al., 2018). Toutes les connaissances et l'information de nature géoscientifique de la Carte préliminaire 7-1981 qui sont en conformité avec le modèle de données ont été conservées pendant le processus de conversion. Le but de la conversion de cartes publiées antérieurement suivant un langage scientifique commun et une légende commune est de permettre et de faciliter la compilation, l'interprétation, la gestion et la diffusion efficaces de l'information géologique cartographique en mode numérique de façon structurée et cohérente. Cette façon de faire offre un outil efficace de gestion des connaissances élaboré à l'aide d'une géodatabase qui pourra évoluer suivant le type d'information à paraître sur les nouvelles cartes de la géologie des formations superficielles.



National Topographic System reference and index to adjoining published Geological Survey of Canada maps

Catalogue No. M183-1/385-2018E-PDF  
ISBN 978-0-660-28175-6  
<https://doi.org/10.4095/313098>

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources, 2019


 Natural Resources Canada  
Ressources naturelles Canada

## CANADIAN GEOSCIENCE MAP 385

### SURFICIAL GEOLOGY

# LOWER HAYES RIVER

Nunavut  
NTS 56-M  
1:250 000



#### QUATERNARY

##### POST LAST GLACIATION

###### NONGLACIAL ENVIRONMENT

**EOLIAN SEDIMENTS:** silt and fine sand, moderately sorted and laminated, in places with cross-stratification, ripple marks, and finely disseminated organic material; less than 2 m thick; deposits too small to be mapped are generally included with units Ap, At, Gf, and GF.

**Eolian sediments, undifferentiated:** silt and fine sand; less than 2 m thick; unvegetated, presently active.

**ALLUVIAL SEDIMENTS:** silt, sand, and fine gravel; moderately to well sorted but commonly interstratified with beds of distinctly different grain sizes; crossbedding, scour-and-fill structures, and ripple marks common; up to 5 m thick.

**Floodplain sediments:** silt, sand, and gravel; variable thickness; seasonally flooded, unvegetated.

**Alluvial terraced sediments:** silt, sand, and gravel; variable thickness; above present flood zone, vegetated.

**Alluvial veneer:** silt, sand, and fine gravel; commonly less than 1 m thick; overlying bedrock.

##### LAST GLACIATION

###### PROGLACIAL AND GLACIAL ENVIRONMENT

**MARINE SEDIMENTS:** silt, sand, and in places clay or gravel; generally less than 10 m but up to 75 m thick; deposited in high stand of the sea during glacial retreat.

**Beach sediments:** sand and silt or gravel, horizontal and cross-stratified; commonly less than 1 m thick; occurs as beaches overlying bedrock.

**Littoral and nearshore sediments:** sand and silt or gravel, horizontal and cross-stratified; from 2 m to a few metres thick; occurs as beaches and terraces; may locally overlie bedrock.

**Offshore sediments:** silt and clay, commonly rhythmically bedded; generally less than 10 m but up to 75 m thick; deposited in deep-water environments.

**Glaciomarine veneer:** silt, sand, and in places clay or gravel; commonly less than 1 m thick; deposited in deep-water environments; overlying bedrock.

**GLACIOFLUVIAL SEDIMENTS:** sand and gravel deposited in, or around, or near a glacier, largely as a result of meltwater stream flow.

**Ice-contact sediments:** sand and gravel, irregular to cross-stratified with poor to moderate sorting; up to 30 m thick; occurs as ice-contact deltas, sinuous ridges, isolated hummocks, local blanket deposits, and esker and kame complexes; may locally overlie bedrock.

**Glaciofluvial sediments, undifferentiated:** sand and gravel, massive to well stratified and sorted; up to 70 m thick; with smooth, flat to inclined surfaces; occurs as deltas, fans, or terraced valley-fill deposits.

###### GLACIAL ENVIRONMENT

**GLACIAL SEDIMENTS (TILL):** poorly sorted till (diamicton) deposited directly by glacial ice.

**Till veneer:** silty, gravelly sand (diamicton) with less than 10% clay, nonsorted, nonstratified; commonly less than 1 m thick overlying bedrock; boulders common on the surface; occurs as veneer on underlying bedrock or as thin hummocks and ridges (moraines); where washed scoured lag overlay is present, the upper metre is abnormally sandy due to either removal of fines by wave action or intermixing of marine, glaciolacustrine, or glaciofluvial sand.


**Till, undifferentiated:** silty, gravelly sand (diamicton) with less than 10% clay, nonsorted, nonstratified, compact but un lithified; generally 7 m up to 20 m thick; boulders up to 2 m long common on the surface; occurs mainly as blanket on underlying bedrock or as hummocks and ridges (moraines); where washed scoured lag overlay is present, the upper metre is abnormally sandy due to either removal of fines by wave action or intermixing of marine, glaciolacustrine, or glaciofluvial sand.

##### PRE-QUATERNARY

**Bedrock, undifferentiated:** Precambrian igneous and metamorphic crystalline rock of variable composition and structure.

**Complex units:** two map-unit designators separated by a dot (.) are used where the surficial cover forms a complex area and the units are too small to be mapped individually (e.g. Tv.GfC designates an area of till veneer with numerous patches of ice-contact sediments).

**Stratigraphic relationship:** two map-unit designators separated by a slash (/) are used where a stratigraphic relationship is observed or confidently inferred (e.g. Tv/R designates till veneer overlying bedrock).



**Recommended citation**  
Geological Survey of Canada, 2019, Surficial geology, lower Hayes River, Nunavut, NTS 56-M; Geological Survey of Canada, Canadian Geoscience Map 385 (Surficial Data Model v. 2.3.14 conversion of Map 7-1981), scale 1:250 000. <https://doi.org/10.4095/313098>

## CANADIAN GEOSCIENCE MAP 385

### SURFICIAL GEOLOGY

# LOWER HAYES RIVER

Nunavut

NTS 56-M

1:250 000



**Author:** Geological Survey of Canada

Geology by R.D. Thomas (east half) and A.S. Dyke (west half), 1976 and 1977

Geological compilation by R.D. Thomas

Geology conforms to Surficial Data Model v. 2.3.14 (Deblonde et al., 2018).

Geological data conversion by D. E. Kerr, 2012, 2017, and 2018

Geomatics by S. Eagles, K. McNeil, C.D. Stevens, and J. Kingsley

Cartography by N. Côté

Scientific editing by A. Weatherston

Initiative of the Geological Survey of Canada, conducted under the auspices of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) Program

Map projection Universal Transverse Mercator, zone 15 North American Datum 1983

Base map at the scale of 1:250 000 from Natural Resources Canada, with modifications  
Elevations in metres above mean sea level

Proximity to the North Magnetic Pole causes the magnetic compass to be erratic in this area.

Mean magnetic declination 2019, 7°27'W, decreasing 4.6' annually  
Readings vary from 9°38'W in the NE corner to 5°21'W in the SW corner of the map.

This map is not to be used for navigational purposes.

The Geological Survey of Canada welcomes corrections or additional information from users.

Data may include additional observations not portrayed on this map. See map info document accompanying the downloaded data for more information about this publication.

This publication is available for free download through GEOSCAN (<https://geoscan.nrcan.gc.ca/>).

## CANADIAN GEOSCIENCE MAP 385

### SURFICIAL GEOLOGY

# LOWER HAYES RIVER

Nunavut

NTS 56-M

