

TABLE 1: AVERAGE ANNUAL LANDED VALUE OF INLAND COMMERCIAL FISHERIES, 1959-1963-LEADING SPECIES

CANADA	\$'000	N.W.T.	ALBERTA	SASKATCHEWAN	MANITOBA	ONTARIO	QUEBEC	N.B.
	\$'000	\$'000	\$'000	\$'000	\$'000	\$'000	\$'000	\$'000
A Lake Whitefish <i>Coregonus clupeaformis</i>	3,612.3	A 534.2	A 509.4	A 734.9	A 955.4	A 863.4	A 15.0 *	
B Walleye <i>Stizostedion vitreum</i>	3,084.2	B 13.9 *	B 86.5	B 282.2	B 1,512.1	B 1,119.9	B 66.6 **	B 3.0 *
C Yellow Perch <i>Perca flavescens</i>	1,520.3		C 7.0 *		C 53.9	C 1,445.8	C 13.6 *	
D Sauger <i>Stizostedion canadense</i>	903.7				D 880.2	D 23.5		
E Tullibee <i>Coregonus artedii</i>	756.9		E 199.6	E 34.4	E 46.3	E 476.6		
F Lake Trout <i>Salvelinus namaycush</i>	557.8	F 173.7	F 7.5 *	F 208.7	F 61.1	F 99.7	F 7.1 *	
G Northern Pike <i>Esox lucius</i>	436.4	G 12.0 *	G 54.7	G 57.4	G 216.5	G 83.4	G 12.4 *	
H Rainbow Smelt <i>Osmerus mordax</i>	406.0				H 397.4	H 8.6 *		
I Lake Sturgeon <i>Acipenser fulvescens</i>	292.6			I 1.6 *	I 19.3	I 156.6	I 113.1	I 2.0 *
J White Bass <i>Morone chrysops</i>	275.2				J 275.2	J 130.7		
K Channel Catfish <i>Ictalurus punctatus</i>	205.6				K 7.2 *	K 19.6	K 67.7	
L American Eel <i>Anguilla rostrata</i>	140.1				L 19.6	L 119.5	L 1.0 *	
M Sucker <i>Catostomus spp</i>	120.7		M 11.6 *		M 83.0	M 26.0	M 0.1 *	
N Carp <i>Cyprinus carpio</i>	100.6				N 2.4 *	N 84.9	N 13.3 *	
O Alewife <i>Alosa pseudoharengus</i>	85.7							O 85.7
P Minnow Family <i>Cyprinidae</i>	49.0						P 49.0	
Q Burbot <i>Lota lota</i>	29.5		Q 11.8 *		Q 14.3 *	Q 3.4 *		
R Atlantic Tomcod <i>Microgadus tomcod</i>	26.1						R 26.1	
TOTAL	12,602.7	733.8	888.1	1,319.2	3,851.7	5,206.1	512.0	91.8

TABLE 2: AVERAGE ANNUAL LANDED VALUE 1959-1963 - MINOR SPECIES

The following inland species plus "Other" collectively had an average landed value for the period 1959-1963 of \$190,000 or 1.4% of the national average value of inland fisheries.

SPECIES	CAN.	N.W.T.	ALTA.	SASK.	MAN.	ONT.	QUE.	N.B.
Atlantic Salmon <i>Salmo salar</i>	\$15,600							15,600
Goideye <i>Hiodon alosoides</i>	\$15,300		400		11,000	3,900		
American Shad <i>Alosa sapidissima</i>	\$14,700						2,900	11,800
Inconnu <i>Stenodus leucithys</i>	\$13,400	13,400						
Freshwater Drum <i>Aplodinotus grunniens</i>	\$13,300				13,300			
Striped Bass <i>Morone saxatilis</i>	\$8,000						1,100	6,900
Crappie <i>Promoxis spp</i>	\$2,800						2,800	
Other	\$107,000	100	700	26,400		77,300	2,500	
TOTAL	\$190,100	13,500	1,100	26,400	24,300	81,200	9,300	34,300

The dots shown in Table 1 by species have the same value as the dots distributed on the maps on pages 170, 171, and 172 and are intended to facilitate visual comparisons of the values of inland commercial fisheries with those of the Atlantic and Pacific fisheries. Each dot represents \$23,082 or .02% of the average annual national landed value of fisheries for the period 1959-1963.

* Less than the value for one dot.

