
SEASONAL RUNOFF
PERIODS OF MAXIMUM AND MINIMUM STREAMFLOW
February
March
January
April
December
May
November
June
October
July
September
August
Coloured discs indicate the month of maximum streamflow at gauging stations. The location of each station is shown by a dot in the center of each disc. Solid dots (•) indicate that the period of minimum streamflow occurs between October and April inclusive. Open dots (•) indicate that minimum flow occurs between May and September inclusive. The period of record at gauging stations is of varying length and continuity.
ALASKA
(UNITED STATES OF AMERICA)
YUKON TERRITORY
NORTHWEST TERRITORIES
GREENLAND
(DENMARK)
BRITISH COLUMBIA
ALBERTA
SASKATCHEWAN
MANITOBA
ONTARIO
QUEBEC
NEWFOUNDLAND
NEW BRUNSWICK
PRINCE EDWARD ISLAND
UNITED STATES OF AMERICA
PACIFIC OCEAN
ATLANTIC OCEAN
Beaufort Sea
Baffin Bay
Davis Strait
Hudson Bay
Hudson Strait
Labrador Sea
Gulf of St. Lawrence
Lake Superior
Lake Michigan
Lake Huron
Lake Ontario
Lake Erie
Great Slave Lake
Great Bear Lake
Lake Athabasca
Lake Winnipeg
Lake Manitoba
Lake Nipigon
Lake Abitibi
Lake Mistassini
Lake Nipissing
Lake Simcoe
Lake St. Clair
Lake Kamouraska
Lake Ste. Anne
Lake Ste. Marie
Lake Ste. Claire