

Atlas of Canada 6th Edition  
 (archival version)

Break-up of Sea Ice

The typical retreat of the sea ice cover from the late winter to late summer is shown on this map. Sea ice is any form of ice that is found at sea and has originated from the freezing of seawater. Melting of sea ice begins in spring in the Gulf of St. Lawrence and East Newfoundland, retreating northward towards the Labrador coast. In June openings appear in Baffin Bay and the Beaufort Sea, while clearing is already underway in Hudson Bay. Break-up continues throughout the summer months, reaching a minimum extent around mid-September. The map shows the typical retreat of the sea ice cover from late winter to late summer (mid-September) on a bi-weekly basis. The dates shown in the legend of the map indicate the extent of the sea ice cover at those dates.


0 150 300 450 600 km

Lambert Conformal Conic Projection. Standard Parallels 49°N and 77°N

<b>Break-up of Sea Ice</b>	<b>Populated Places</b>	<b>Road network</b>
March 15	• 1-4 999	— Road network
April 1	• 5 000-49 999	— Ferry route
April 15	• 50 000-99 999	<b>Boundaries</b>
May 1	• 100 000 and greater	— International
May 15	★ Provincial and Territorial Capital	— Provincial / Territorial
June 1		— EEZ (200 mile)
June 15		— Canada / Kaiaallit Nunaat dividing line
July 1		
July 15		
August 1		
August 15		
Sept. 01 to Oct 01 (irregular breakup)		
Permanent ice		

**Source(s):**  
 Break-up of Sea Ice  
 Regional Ice Charts from 1969 to 1998, Canadian Ice Service, Environment Canada.

© 2009, Her Majesty the Queen in Right of Canada, Natural Resources Canada.



This product was originally published as an interactive map in the online Atlas of Canada, 6th Edition. It was not intended for high resolution reproduction and therefore does not conform to the standards of cartographic representation and map design found on the Atlas of Canada's paper reference maps.